arXiv:1003.1317v1 [math.RT] 5 Mar 2010

REPRESENTATIONS OF THREAD QUIVERS

CARL FREDRIK BERG AND ADAM-CHRISTIAAN VAN ROOSMALEN

Abstract. We characterize the small k-linear (k algebraically closed) Karoubi categories of which the category of finitely presented representations are abelian and satisfy Serre duality. Furthermore, we show that those categories whose category of finitely presented representations is hereditary are classified by so called thread quivers.

Contents

1. Introduction 1
2. Preliminaries 3
2.1. Notations and conventions 3
2.2. Serre duality 4
2.3. 2-categories and 2-colimits 4
2.4. Free categories 5
2.5. Representations of preadditive categories 6
3. Preadditive semi-hereditary categories 7
4. Dualizing k-varieties 7
5. (Co)reflective subvarieties 11
6. Threads 13
7. Classification by thread quivers 15
8. Representations of thread quivers 21
Appendix A. 2-Pushouts 22
References 25

1. Introduction

It was shown in [6] that every k-linear (where k is an algebraically closed field) abelian noetherian hereditary category with Serre duality which is generated by preprojective objects is derived equivalent to the category $\operatorname{rep} Q$ of finitely presented representations of a strongly locally finite quiver (i.e. a quiver where the indecomposable projective and injective representations have finite length).

In [5] we wish to remove the noetherian condition. It was shown in [15] that in this more general case, the abelian categories under consideration cannot be derived equivalent to a noetherian category such as $\operatorname{rep} Q$. In order to accommodate these larger nonnoetherian categories, we need to generalize our notion of a quiver to a suitable category $\mathfrak{a}$, where "suitable" means that the category $\operatorname{mod} \mathfrak{a}$ of finitely presented modules is abelian, hereditary, and has Serre duality. It turns out that these categories can be classified by thread quivers (see Theorem 1.2 below).

In this way, we contribute to the study of dualizing k-varieties (introduced in [2]) and we expand upon the results of [16].

Before stating our results, we will give some definitions. A *finite k-variety* is a Hom-finite additive category $\mathfrak{a}$ where idempotents split. The functors $\mathfrak{a}(-, A)$ and $\mathfrak{a}(A, -)^*$ from $\mathfrak{a}$ to $\operatorname{mod} k$ will be called *standard projective representations* and *standard injective representations*, respectively. We will write $\operatorname{mod} \mathfrak{a}$ for the category of contravariant functors $\mathfrak{a} \to \operatorname{mod} k$ which are finitely presentable by standard projectives.

$$\begin{array}{ccccc} \mathbb{N} & \cdot & \mathcal{P}_t \overset{\rightarrow}{\times} \mathbb{Z} & \cdot & -\mathbb{N} \\ \boxed{0 \to 1 \to 2 \to \cdots} & \cdots & \boxed{\cdots \to (p,-1) \to (p,0) \to (p,1) \to \cdots} & \cdots & \boxed{\cdots \to -2 \to -1 \to -0} \end{array}$$

FIGURE 1. The poset $\mathcal{L}_t$

We will say a finite k-variety $\mathfrak{a}$ is *dualizing* if and only if (Proposition 4.1) $\mathfrak{a}$ has pseudokernels and pseudocokernels (thus $\operatorname{mod} \mathfrak{a}$ and $\operatorname{mod} \mathfrak{a}^\circ$ are abelian, where $\mathfrak{a}^\circ$ is the dual category of $\mathfrak{a}$), every standard projective object is cofinitely generated by standard injectives, and every standard injective object is finitely generated by standard projectives.

A finite k-variety $\mathfrak{a}$ is called *semi-hereditary* if and only if the category $\operatorname{mod} \mathfrak{a}$ is abelian and hereditary. It has been shown (see Proposition 3.1) that $\mathfrak{a}$ is semi-hereditary if and only if every full (preadditive) subcategory with finitely many elements is semi-hereditary.

Let $\mathfrak{a}$ be a finite k-variety. The following theorem (Theorem 4.7) shows that $\operatorname{mod} aa$ is an abelian and hereditary category with Serre duality if and only if $\mathfrak{a}$ is a semi-hereditary dualizing (finite) k-varieties.

Theorem 1.1. *Let $\mathfrak{a}$ be a finite k-variety. The following are equivalent:*

1. $\operatorname{mod} \mathfrak{a}$ *is abelian and has Serre duality,*
2. $\mathfrak{a}$ *is a dualizing k-variety and every object of* $\operatorname{mod} \mathfrak{a}$ *has finite projective and finite injective dimension.*

Intuitively, a semi-hereditary dualizing k-variety can be seen as the free k-linear path category kQ of a strongly locally finite quiver k without relations, but where some arrows are replaced by infinite locally discrete (= without accumulation points) linearly ordered posets. To accommodate this extra information, we will introduce thread quivers.

A *thread quiver* consists of the following information:

- A quiver $Q = (Q_0, Q_1)$ where Q_0 is the set of vertices and Q_1 is the set of arrows.
- A decomposition $Q_1 = Q_s \coprod Q_t$. Arrows in Q_s will be called *standard arrows*, while arrows in Q_t will be referred to as *thread arrows*.
- With every thread arrow t, there is an associated linearly ordered set $\mathcal{P}_t$, possibly empty.

With every thread arrow $t : x \overset{\mathcal{P}_t}{\dashrightarrow} y$ in Q we associate a locally discrete (= there are no accumulation points) linearly ordered poset $\mathcal{L}_t$ with a minimal and a maximal element: namely $\mathcal{L}_t = \mathbb{N} \cdot (\mathcal{P}_t \overset{\rightarrow}{\times} \mathbb{Z}) \cdot -\mathbb{N}$, where $A \cdot B$ means "first A, then B" and $\mathcal{P}_t \overset{\rightarrow}{\times} \mathbb{Z}$ is the poset $\mathcal{P}_t \times \mathbb{Z}$ with the lexicographical ordering (see Figure 1).

The poset $\mathcal{L}_t$ is interpreted as a category in the usual sense, and $k\mathcal{L}_t$ will denote the k-linearized category.

To recuperate the semi-hereditary dualizing k-variety from the thread quiver Q, we need to replace the thread arrows by the corresponding linearly ordered posets. Since is cumbersome –albeit possible– to do this "by hand," we will prefer a more global approach: by a 2-pushout.

For a thread quiver Q, denote by Q_r the underlying "regular" quiver, thus where all the thread arrows are regular (and unlabeled) arrows. By kQ_r we will denote the normal k-linear additive path category of Q_r.

With every thread arrow $t : x_t \overset{\mathcal{P}_t}{\dashrightarrow} y_t$ of Q, there is an associated functor $k(x_t \to y_t) \to kQ_r$ and a functor $k(x_t \to y_t) \longrightarrow k\mathcal{L}_t$ where $\mathcal{L}_t = \mathbb{N} \cdot (\mathcal{P}_t \overset{\rightarrow}{\times} \mathbb{Z}) \cdot -\mathbb{N}$ as above. The semi-hereditary dualizing k-variety kQ is then defined as a 2-pushout of

$$\begin{tikzcd} \bigoplus_{t \in Q_t} k(\cdot \to \cdot) \arrow[r, "f"] \arrow[d, "g"'] & kQ_r \\ \bigoplus_{t \in Q_t} k\mathcal{L}_t & \end{tikzcd}$$

effectively replacing the thread arrows in Q by the required linearly ordered posets. The classification of semi-hereditary dualizing k-varieties is then given by the following theorem (Theorem 7.18 in the text). A thread quiver Q will be called strongly locally finite if and only if the underlying quiver Q_r is strongly locally finite (i.e. the indecomposable projective and injective representations have finite dimension as k-vector spaces).

Theorem 1.2. *Every semi-hereditary dualizing k-variety is equivalent to a category kQ where Q is a strongly locally finite thread quiver.*

An added advantage of the construction of kQ as a 2-pushout is that we gain the following description of the category $\operatorname{rep}_k Q = \operatorname{mod} kQ$: the objects of the category $\operatorname{rep} Q$ are given by the following data

1. a finitely presented representation $N(-) : kQ_r \to \operatorname{mod} k$ of kQ_r,
2. for every thread t, a finitely presented representation $L_t(-) : \mathcal{L}_t \to \operatorname{mod} k$, and
3. a natural equivalence $\alpha : \oplus_t L_t(g-) \Rightarrow N(f-)$.

The morphisms are given by the modifications, thus given the data $(N, \{L_t\}_t, \alpha)$ and $(N', \{L'_t\}_t, \alpha')$ of two representations, a morphism is given by

1. a natural transformation $\beta : N \Rightarrow N'$,
2. a natural transformation $\gamma : \oplus_t L_t \Rightarrow \oplus_t L'_t$

such that the following diagram commutes.

$$\begin{tikzcd}
\oplus_t L_t(g-) \arrow[r, "\alpha"] \arrow[d, "\gamma\bullet 1_g"'] & N(f-) \arrow[d, "\beta\bullet 1_f"] \\
\oplus_t L'_t(g-) \arrow[r, "\alpha'"'] & N'(f-)
\end{tikzcd}$$

Acknowledgments The authors like to thank Idun Reiten, Sverre Smalø, and Michel Van den Bergh for many useful discussions and helpful ideas. We would also like to thank Tobias Fritz for pointing us to [18]. The second author gratefully acknowledges the hospitality of the Max-Planck-Institut für Mathematik in Bonn and the Norwegian University of Science and Technology.

2. Preliminaries

2.1. **Notations and conventions.** Throughout, we will always assume that k is an algebraically closed field. All considered categories will be assumed to be k-linear unless explicitly mentioned. We will fix a Grothendieck universe $\mathcal{U}$. All categories are assumed to be $\mathcal{U}$-categories, i.e. all Hom-sets lie in $\mathcal{U}$. We will say that a category is small when the objects form a set in $\mathcal{U}$, and is skeletally small when the isomorphism classes form a set in $\mathcal{U}$.

When $\mathcal{P}_1, \leq_1$ and $\mathcal{P}_2, \leq_2$ are posets, we will denote by $\mathcal{P}_1 \cdot \mathcal{P}_2, \leq$ the poset with underlying set $\mathcal{P}_1 \coprod \mathcal{P}_2$ and with a partial ordering $\leq$ given by

$$a \leq b \Leftrightarrow \begin{cases} a, b \in \mathcal{P}_1 \text{ and } a \leq_1 b \\ a, b \in \mathcal{P}_2 \text{ and } a \leq_2 b \\ a \in \mathcal{P}_1 \text{ and } b \in \mathcal{P}_2 \end{cases}$$

If $\mathcal{P}_1, \leq_1$ and $\mathcal{P}_2, \leq_2$ are posets, we will write $\mathcal{P}_1 \overset{\rightarrow}{\times} \mathcal{P}_2, \leq$ for the partially ordered set with underlying set $\mathcal{P}_1 \times \mathcal{P}_2$ endowed with the lexicographical ordering.

When writing a poset we will usually repress the partial ordering, thus writing $\mathcal{P}$ instead of $\mathcal{P}, \leq$. We will always interpret a poset as a small category in the usual way.

Let $\mathcal{C}$ be a Krull-Schmidt category. By $\operatorname{ind} \mathcal{C}$ we will denote a set of chosen representatives of isomorphism classes of indecomposable objects of $\mathcal{C}$. If $\mathcal{C}'$ is a Krull-Schmidt subcategory of $\mathcal{C}$, we will assume $\operatorname{ind} \mathcal{C}' \subseteq \operatorname{ind} \mathcal{C}$.

For $X, Y \in \operatorname{ind} \mathcal{C}$, we will denote $[X, Y]$ for the full replete (=closed under isomorphisms) additive subcategory of $\mathcal{C}$ such that for all $Z \in \operatorname{ind} \mathcal{C}$, we have $Z \in \operatorname{ind}[X, Y] \Leftrightarrow \mathcal{C}(X, Z) \neq 0$ and $\mathcal{C}(Z, Y) \neq 0$. The subcategories $[X, Y[,]X, Y]$ and $]X, Y[$ are defined in a straightforward manner.

2.2. **Serre duality.** Let $\mathcal{C}$ be a Hom-finite triangulated k-linear category. A *Serre functor* [7] on $\mathcal{C}$ is an additive auto-equivalence $\mathbb{S} : \mathcal{C} \to \mathcal{C}$ such that for every $X, Y \in \operatorname{Ob} \mathcal{C}$ there are isomorphisms

$$\operatorname{Hom}(X, Y) \cong \operatorname{Hom}(Y, \mathbb{S}X)^*$$

natural in X and Y, and where $(-)^*$ is the vector-space dual.

We will say $\mathcal{C}$ has *Serre duality* if $\mathcal{C}$ admits a Serre functor. An abelian category $\mathcal{A}$ is said to satisfy Serre duality when the bounded derived category $D^b\mathcal{A}$ has Serre duality.

2.3. **2-categories and 2-colimits.** We will give a brief overview of the notation and terminology of 2-categories we will use. Our main references are [8, 18]. Let $\mathfrak{Cat}$ be the 1-category of all small categories, thus the objects are given by small categories and the morphisms by functors.

A *(strict) 2-category* is a category enriched over $\mathfrak{Cat}$, also called a $\mathfrak{Cat}$-category (see [9, 13]). Our main example will be $k\mathfrak{AdCat}$, the 2-category of all small k-linear additive categories:

- the 0-cells are given by small k-linear additive categories,
- the 1-cells are the k-linear functors,
- the 2-cells are natural transformations.

Composition of 1-cells are denoted by $\circ$. Following [14, 18] we will write $\circ$ for vertical composition of 2-cells and $\bullet$ for horizontal composition. We have the following equation

$$(\gamma \bullet \delta) \circ (\alpha \bullet \beta) = (\gamma \circ \alpha) \bullet (\delta \circ \beta) \tag{1}$$

We proceed to define a 2-colimit over a 2-functor.

Definition 2.1. Let $\mathcal{I}$ be a small 1-category. A *(strict) 2-functor* $\mathfrak{a} : \mathcal{I} \to k\mathfrak{AdCat}$ is given by the following data:

(1) a 0-cell $\mathfrak{a}(i)$ of $k\mathfrak{AdCat}$ for every $i \in \operatorname{Ob} \mathcal{I}$, and
(2) a 1-cell $\mathfrak{a}(s) : \mathfrak{a}(i) \to \mathfrak{a}(j)$ of $k\mathfrak{Cat}$ for every morphism $s : i \to j$ in $\mathcal{I}$.

such that $\mathfrak{a}(1_i) = 1_{\mathfrak{a}(i)}$ for all $i \in \operatorname{Ob} \mathcal{I}$ and $\mathfrak{a}(t \circ s) = \mathfrak{a}(t) \circ \mathfrak{a}(s)$ for all composable morphisms $s, t \in \operatorname{Ob} \mathcal{I}$.

Remark 2.2. A (strict) 2-functor is just a functor from $\mathcal{I}$ to the underlying 1-category of $k\mathfrak{AdCat}$.

Example 2.3. For every object $\mathcal{C}$ of $k\mathfrak{AdCat}$, there is a 2-functor $\mathcal{C} : \mathcal{I} \to k\mathfrak{AdCat}$ sending every object of $\mathcal{I}$ to $\mathcal{C}$ and sending every morphism of $\mathcal{C}$ to the identity on $\mathcal{C}$.

Definition 2.4. Let $\mathfrak{a}, \mathfrak{b} : \mathcal{I} \to k\mathfrak{AdCat}$ be two 2-functors. A 2-natural transformation $\mathfrak{f} : \mathfrak{a} \to \mathfrak{b}$ between 2 diagrams consists of the following data:

(1) a 1-cell $\mathfrak{f}_i : \mathfrak{a}(i) \to \mathfrak{b}(i)$ of $k\mathfrak{AdCat}$ for every $i \in \operatorname{Ob} \mathcal{I}$, and
(2) a natural equivalence $\theta^{\mathfrak{f}}_s : \mathfrak{b}(s) \circ \mathfrak{f}_i \to \mathfrak{f}_j \circ \mathfrak{a}(s)$ for every morphism $s : i \to j$ in $\mathcal{I}$.

such that for any two composable morphisms $s : i \to j$, $t : j \to k$ in $\mathcal{I}$, we have the following commutative diagram

$$\begin{tikzcd}
\mathfrak{b}(t \circ s) \circ \mathfrak{f}_i \arrow[r, equal] \arrow[dd, "\theta^{\mathfrak{f}}_{t \circ s}"'] & \mathfrak{b}(t) \circ \mathfrak{b}(s) \circ \mathfrak{f}_i \arrow[d, "1_{\mathfrak{b}(t)} \bullet \theta^{\mathfrak{f}}_s"] \\
& \mathfrak{b}(t) \circ \mathfrak{f}_i \circ \mathfrak{a}(s) \arrow[d, "\theta^{\mathfrak{f}}_t \bullet 1_{\mathfrak{a}(s)}"] \\
\mathfrak{f}_k \circ \mathfrak{a}(t \circ s) \arrow[r, equal] & \mathfrak{f}_k \circ \mathfrak{a}(t) \circ \mathfrak{a}(s)
\end{tikzcd}$$

Definition 2.5. Let $\mathfrak{a}, \mathfrak{b} : \mathcal{I} \to k\mathfrak{AdCat}$ be two 2-functors and $\mathfrak{f}, \mathfrak{g} : \mathfrak{a} \to \mathfrak{b}$ be 2-natural transformations. A *modification* $\Lambda : \mathfrak{f} \to \mathfrak{g}$ consists in giving a 2-cell $\Lambda_i : \mathfrak{f}_i \to \mathfrak{g}_i$ for all objects $i \in \mathcal{I}$ such that for all $s : i \to j$ in $\mathcal{I}$ the following diagram commutes

$$\begin{tikzcd}
\mathfrak{b}(s) \circ \mathfrak{f}_i \arrow[r, "\theta^{\mathfrak{f}}_s"] \arrow[d, "1_{\mathfrak{b}(s)} \bullet \Lambda_i"'] & \mathfrak{f}_j \circ \mathfrak{a}(s) \arrow[d, "\Lambda_j \bullet 1_{\mathfrak{a}(s)}"] \\
\mathfrak{b}(s) \circ \mathfrak{g}_i \arrow[r, "\theta^{\mathfrak{g}}_s"] & \mathfrak{g}_j \circ \mathfrak{a}(s)
\end{tikzcd}$$

Definition 2.6. The diagrams, 2-natural transformations, and modifications form a (strict) 2-category called $2\mathcal{F}(\mathcal{I}, k\mathfrak{AdCat})$.

We can now give the definition of a 2-colimit.

Definition 2.7. Let $\mathfrak{a} : \mathcal{I} \to k\mathfrak{AdCat}$ be a 2-functor. We say $\mathfrak{a}$ admits a 2-colimit if and only if there exist

(1) a k-linear additive category $2\operatorname{colim}\mathfrak{a}$, and
(2) a 2-natural transformation $\sigma : \mathfrak{a} \to 2\operatorname{colim}\mathfrak{a}$,

such that for every category $\mathcal{C}$ the functor

$$(-\circ\sigma) : \operatorname{Hom}_{k\mathfrak{AdCat}}(2\operatorname{colim}\mathfrak{a}, \mathcal{C}) \to \operatorname{Hom}_{2\mathcal{F}}(\mathfrak{a}, \mathcal{C})$$

is an equivalence of categories.

Remark 2.8. The fact that $(-\circ\sigma)$ is fully faithful can be restated as follows: Let $\mathfrak{f}, \mathfrak{g} : \mathfrak{a} \to \mathcal{C}$ be 2-natural transformation and let $\Lambda : \mathfrak{f} \to \mathfrak{g}$ be a modification. Let $f, g : 2\operatorname{colim}\mathfrak{a} \to \mathcal{C}$ be two functors corresponding with $\mathfrak{f}$ and $\mathfrak{g}$, respectively. Then there is a unique natural transformation $\alpha : f \to g$ such that the following diagram commutes.

$$\begin{tikzcd}
\mathfrak{f}_i \arrow[r, "\Lambda_i"] \arrow[d] & \mathfrak{g}_i \arrow[d] \\
f\circ\sigma_i \arrow[r, "\alpha\bullet 1_{\sigma_i}"'] & g\circ\sigma_i
\end{tikzcd}$$

We have the following result ([18, Theorem A.3.4]).

Theorem 2.9. *Denote by* $\mathfrak{Cat}$ *the 2-category of all small categories. Let* $\mathcal{I}$ *be a small 1-category, and* $\mathfrak{a} : \mathcal{I} \to \mathfrak{Cat}$ *a 2-functor. Then* $\mathfrak{a}$ *admits a 2-colimit.*

One may construct this 2-colimit as follows. We start by constructing a category $\int_{\mathcal{I}}\mathfrak{a}$. We set

$$\operatorname{Ob}\int_{\mathcal{I}}\mathfrak{a} = \{(i, X) \mid i \in \operatorname{Ob}\mathcal{I}, X \in \operatorname{Ob}\mathfrak{a}(i)\} = \coprod_{i\in\mathcal{I}} \operatorname{Ob}\mathfrak{a}(i).$$

and

$$\operatorname{Hom}((i, X), (j, Y)) = \{(s, f) \mid s \in \mathcal{I}(s, t), f \in \operatorname{Hom}_{\mathfrak{a}(j)}(\mathfrak{a}(s)(X), Y)\}.$$

The composition is defined as $(t, g) \circ (s, f) = (t \circ s, g \circ \mathfrak{a}(t)(f))$. Note that for every $i \in \operatorname{Ob}\mathcal{I}$ there is a natural functor $\mathfrak{a}(i) \to \int_{\mathcal{I}}\mathfrak{a}$.

We consider the set SS of morphisms of $\int_{\mathcal{I}}\mathfrak{a}$ given by

$$SS = \{(s, f) : (i, X) \to (j, Y) \mid f : \mathfrak{a}(s)(X) \to Y \text{ is an isomorphism}\}.$$

It is then shown that $\left(\int_{\mathcal{I}}\mathfrak{a}\right)[SS^{-1}]$ together with the canonical functors $\mathfrak{a}(i) \to \left(\int_{\mathcal{I}}\mathfrak{a}\right)[SS^{-1}]$ is a 2-colimit of $\mathfrak{a}$.

2.4. **Free categories.** A quiver Q is a pair (Q_0, Q_1) consisting of a small set Q_0 of vertices and a small set Q_1 of arrows between the vertices such that between two elements of Q_0, the set of arrows is a small set. Note that we allow multiple arrows and loops. A map $f : Q \to Q'$ between quivers consists of a map $f_0 : Q_0 \to Q'_0$ of objects and a map $f_1 : Q_1 \to Q'_1$ of arrows such that $f_1(A \to B)$ is an arrow from $f_0(A)$ to $f_0(B)$.

We will say a quiver is *finite* if both Q_0 and Q_1 are finite. A quiver Q is *locally finite* if every vertex is incident to only a finite number of arrows and Q is said to be *strongly locally finite* if it is locally finite and has no infinite paths of the form $\bullet \to \bullet \to \cdots$ or $\cdots \to \bullet \to \bullet$.

Equivalently, a quiver is strongly locally finite if and only if all indecomposable projective and injective representations have finite length.

A commutativity condition on a quiver is a pair of oriented paths, both starting in the same vertex and ending in the same vertex.

We write $\mathfrak{CondQuiver}$ for the category whose objects are given by quivers with a set of commutativity conditions and whose morphisms of are given by morphisms of quivers $Q \to Q'$ mapping a commutativity condition of Q to a commutativity condition of Q'.

Let $\mathfrak{Cat}$ be the strict 1-category of small categories and functors. There is an obvious (faithful) forgetful functor $\mathfrak{Cat} \to \mathfrak{CondQuiver}$ admitting a left adjoint $\operatorname{Free} : \mathfrak{CondQuiver} \to \mathfrak{Cat}$ (see [8, Proposition 5.1.6]). For a quiver Q, the category $\operatorname{Free} Q$ is called the *free category* or the *path category* of Q.

Let $k\mathfrak{Cat}$ be the 2-category of k-linear categories, k-linear functors, and k-linear natural transformations. There is an obvious forgetful 2-functor $k\mathfrak{Cat} \to \mathfrak{Cat}$ forgetting the k-linear structure. This functor admits a left adjoint $\mathfrak{Cat} \to k\mathfrak{Cat}$ ([9, 13]).

The embedding 2-functor $k\mathfrak{AdCat} \to k\mathfrak{Cat}$ from the 2-category of small k-linear additive categories to the 2-category of small k-linear categories also has a left adjoint $k\mathfrak{Cat} \to k\mathfrak{AdCat}$ given by formally adding biproducts.

Thus the forgetful 1-functor $k\mathfrak{AdCat} \to \mathfrak{CondQuiver}$ has a left adjoint. With a slight abuse of notation, we will write this left adjoint as $k : \mathfrak{CondQuiver} \to k\mathfrak{AdCat}$, thus if Q is a quiver, kQ is a k-linear additive category.

2.5. **Representations of preadditive categories.** Let $\mathfrak{a}$ be a small preadditive category. A right $\mathfrak{a}$-module is a contravariant functor from $\mathfrak{a}$ to $\operatorname{Mod} k$, the category of all vector spaces. The category of all right $\mathfrak{a}$ modules is denoted by $\operatorname{Mod} \mathfrak{a}$.

If $f : \mathfrak{a} \to \mathfrak{b}$ is a functor between small pre-additive categories then there is an obvious restriction functor

$$(-)_{\mathfrak{a}} : \operatorname{Mod}(\mathfrak{b}) \to \operatorname{Mod}(\mathfrak{a})$$

which sends N to $N \circ f$. This restriction functor has a left adjoint

$$\mathfrak{b} \otimes_{\mathfrak{a}} - : \operatorname{Mod}(\mathfrak{a}) \to \operatorname{Mod}(\mathfrak{b})$$

which is the right exact functor which sends the projective generators $\mathfrak{a}(-, A)$ in $\operatorname{Mod}(\mathfrak{a})$ to $\mathfrak{b}(-, f(A))$ in $\operatorname{Mod}(\mathfrak{b})$. As usual if f is fully faithful we have $(\mathfrak{b} \otimes_{\mathfrak{a}} N)_{\mathfrak{a}} = N$.

Let M be in $\operatorname{Mod}(\mathfrak{a})$. We will say that M is *finitely generated* if M is a quotient of finitely generated projectives. We say that M is *finitely presented* if M has a presentation

$$P \to Q \to M \to 0$$

where P, Q are finitely generated projectives. It is easy to see that these notions coincides with the ordinary categorical ones. The full subcategory $\operatorname{Mod} \mathfrak{a}$ spanned by the finitely presented modules will be denoted by $\operatorname{mod} \mathfrak{a}$. If $\operatorname{mod} \mathfrak{a}$ is an abelian category, we will say $\mathfrak{a}$ is *coherent.*

Dually we will say that M is *cofinitely generated* if it is contained in a cofinitely generated injective. *Cofinitely presented* is defined in a similar way.

The categorical interpretation of the latter notions is somewhat less clear. However if $\mathfrak{a}$ is Hom-finite then both finitely and cofinitely presented representations correspond to each other under duality (exchanging $\mathfrak{a}$ and $\mathfrak{a}^\circ$).

It is well-known that $\mathfrak{a}$ is coherent if and only if it has pseudokernels, i.e. if for every morphism $A \to B$ in $\mathfrak{a}$ there is a morphism $K \to A$ such that

$$\mathfrak{a}(-, K) \to \mathfrak{a}(-, A) \to \mathfrak{a}(-, B)$$

is exact. Pseudocokernels are defined in a dual way.

With every object A of $\mathfrak{a}$, we may associate a *standard projective* $\mathfrak{a}(-, A)$ and a *standard injective* $\mathfrak{a}(A, -)^*$. It is clear that every finitely generated projective is a direct summand of a standard projective. If $\mathfrak{a}$ has finite direct sums and idempotents split in $\mathfrak{a}$, then every finitely generated projective is isomorphic to a standard projective. Dual notions hold for injective objects.

We will say a map $A \to B$ in $\mathfrak{a}$ is a *radical morphism* if there are no maps $X \to A$ and $B \to X$ such that the composition

$$X \to A \to B \to X$$

is an isomorphism, where X is not zero. The set of all radical maps in $\mathfrak{a}(A, B)$ is denoted by $\operatorname{rad}(A, B)$; it is a subspace of $\mathfrak{a}(A, B)$.

With an indecomposable object $A \in \operatorname{ind} \mathfrak{a}$, we may associate in a straightforward way the *standard simple* object S_A as $\mathfrak{a}(-,A)/\operatorname{rad}(-,A)$.

3. Preadditive semi-hereditary categories

We say that a small preadditive category $\mathfrak{a}$ is *semi-hereditary* if the finitely presented objects $\operatorname{mod}(\mathfrak{a})$ in $\operatorname{Mod}(\mathfrak{a})$ form an abelian and hereditary category. The following proposition ([17], see also [3, Theorem 1.6]) makes it easy to recognize semi-hereditary categories in examples.

Proposition 3.1. *Let $\mathfrak{a}$ be a small preadditive category, then $\mathfrak{a}$ is semi-hereditary if and only if any full subcategory of $\mathfrak{a}$ with a finite number of objects is semi-hereditary.*

Lemma 3.2. *Let $\mathfrak{b} \to \mathfrak{c}$ be a full embedding of preadditive categories. Then $\mathfrak{c} \otimes_{\mathfrak{b}} - : \operatorname{mod}(\mathfrak{b}) \to \operatorname{mod}(\mathfrak{c})$ is fully faithful.*

Proof. This may be checked on objects of the form $\mathfrak{b}(-,B)$ where it is clear. □

Since $(\mathfrak{c} \otimes_{\mathfrak{b}} -, (-)_{\mathfrak{b}})$ is an adjoint pair, if follows from this lemma that $(\mathfrak{c} \otimes_{\mathfrak{b}} -)_{\mathfrak{b}}$ is naturally isomorphic to the identity functor on $\operatorname{mod}(\mathfrak{b})$.

The following result is a slight generalization of [3, Proposition 2.2].

Proposition 3.3. *Let $\mathfrak{b} \to \mathfrak{c}$ be a full embedding of semi-hereditary categories. Then the (fully faithful) functor $\mathfrak{c} \otimes_{\mathfrak{b}} - : \operatorname{mod}(\mathfrak{b}) \to \operatorname{mod}(\mathfrak{c})$ is exact.*

Proof. Let $M \in \operatorname{mod}(\mathfrak{b})$ and consider a projective resolution $0 \to P_1 \xrightarrow{\theta} P_0 \to M \to 0$ in $\operatorname{mod}(\mathfrak{b})$. Then P_1 is a direct summand of some $\oplus_{i=1}^{n} \mathfrak{b}(-,B_i)$. Put $K = \ker(\mathfrak{c} \otimes_{\mathfrak{b}} \theta)$. Then K is a direct summand of $\mathfrak{c} \otimes_{\mathfrak{b}} P_1$ (since $\mathfrak{c}$ is semi-hereditary) and $K_{\mathfrak{b}} = 0$ (since θ is injective).

Assume K is non-zero. Since K is a direct summand of $\mathfrak{c} \otimes_{\mathfrak{b}} P_1$ we obtain a non-zero map $\mathfrak{c} \otimes_{\mathfrak{b}} P_1 \to K$ and hence a non-zero map $\oplus_{i=1}^{n} \mathfrak{c}(-,B_i) \to K$ and thus ultimately a non-zero element of some $K(B_i)$, contradicting the fact that $K_{\mathfrak{b}} = 0$.

Thus $K = 0$. If we denote the left satellites of $\mathfrak{c} \otimes_{\mathfrak{b}} -$ by $\operatorname{Tor}_i^{\mathfrak{b}}(\mathfrak{c}, -)$ then we have just shown that $\operatorname{Tor}_i^{\mathfrak{b}}(\mathfrak{c}, -) = 0$ for $i > 0$. Hence $\mathfrak{c} \otimes_{\mathfrak{b}} -$ is exact. □

4. Dualizing k-varieties

We recall some definitions from [1, 2]. A Hom-finite additive k-linear category where idempotents split is called a *finite k-variety*. Such a finite k-variety is always Krull-Schmidt.

Denote by $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$ the abelian category of locally finite dimensional right $\mathfrak{a}$-modules, thus the full subcategory of $\operatorname{Mod} \mathfrak{a}$ spanned by all contravariant functors from $\mathfrak{a}$ to $\operatorname{mod} k$. Note that an additive k-linear category where idempotents split is a finite k-variety if and only if every standard projective and standard injective lies in $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$.

Let $\mathfrak{a}$ be a finite k-variety. There is a duality $D : \operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a} \to \operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}^\circ$ given by sending a module $M \in \operatorname{Ob} \operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$ to the dual $D(M)$ where $D(M)(x) = \operatorname{Hom}_k(M(x), k)$ for all $x \in \mathfrak{a}$.

If this functor induces a duality $D : \operatorname{mod} \mathfrak{a} \to \operatorname{mod} \mathfrak{a}^\circ$ by restricting to the finitely presented objects in $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$, then we will say that $\mathfrak{a}$ is a *dualizing k-variety*.

The following theorem is a reformulation of [2, Theorem 2.4].

Proposition 4.1. *Let $\mathfrak{a}$ be a finite k-variety, then $\mathfrak{a}$ is a dualizing k-variety if and only if*

1. *$\mathfrak{a}$ has pseudokernels and pseudocokernels, and*
2. *every standard projective is cofinitely presented and every standard injective is finitely presented.*

Proof. Assume $\mathfrak{a}$ is a dualizing k-variety, so that there is a duality $D : \operatorname{mod} \mathfrak{a} \to \operatorname{mod} \mathfrak{a}^\circ$. Since $\operatorname{mod} \mathfrak{a}^\circ$ is a right exact subcategory of $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}^\circ$, we know that $\operatorname{mod} \mathfrak{a}$ is a left exact subcategory of $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$. Hence $\operatorname{mod} \mathfrak{a}$ is closed under kernels and we deduce that $\mathfrak{a}$ has pseudokernels. Likewise, one shows that $\mathfrak{a}^\circ$ has pseudokernels or, dually, that $\mathfrak{a}$ has pseudocokernels.

Let I be a standard injective in $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$, so that DI is a standard projective in $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}^\circ$. We then know that $DI \in \operatorname{Ob} \operatorname{mod} \mathfrak{a}^\circ$ and thus by duality $I \in \operatorname{Ob} \operatorname{mod} \mathfrak{a}$ so that I is finitely presented.

Let P be a standard projective in $\operatorname{mod}\mathfrak{a}$. Since D induces a duality between $\operatorname{mod}\mathfrak{a}$ and $\operatorname{mod}\mathfrak{a}^\circ$, we know that DP is an object in $\operatorname{mod}\mathfrak{a}^\circ$, hence finitely presented. Taking the vector space dual of a projective resolution in $\operatorname{mod}\mathfrak{a}^\circ$ yields an injective resolution in $\operatorname{mod}\mathfrak{a}$. We obtain that P is cofinitely presented.

For the other direction, let $X \in \operatorname{mod}\mathfrak{a}^\circ$. Since X is finitely presented, the dual $DX \in \operatorname{Mod}_{\mathrm{lfd}}\mathfrak{a}$ is cofinitely presented, thus there is an exact sequence

$$0 \to DX \to I \xrightarrow{f} J$$

in $\operatorname{Mod}_{\mathrm{lfd}}\mathfrak{a}$. By assumption $\mathfrak{a}$ has pseudokernels such that $\operatorname{mod}\mathfrak{a}$ is an exact subcategory of $\operatorname{Mod}_{\mathrm{lfd}}\mathfrak{a}$. It now follows that $DX \in \operatorname{mod}\mathfrak{a}$ so that the functor $D : \operatorname{Mod}_{\mathrm{lfd}}\mathfrak{a}^\circ \to \operatorname{Mod}_{\mathrm{lfd}}\mathfrak{a}$ restricts to a functor $D : \operatorname{mod}\mathfrak{a}^\circ \to \operatorname{mod}\mathfrak{a}$.

Similarly, one shows that there is a functor $D : \operatorname{mod}\mathfrak{a} \to \operatorname{mod}\mathfrak{a}^\circ$ and since $D^2 \cong 1$, these are equivalences. This shows that $\mathfrak{a}$ is a dualizing k-variety. □

Remark 4.2. It follows from the proof of Proposition 4.1 that every object in the abelian category $\operatorname{mod}\mathfrak{a}$ has a resolution by objects of $\mathcal{P}$ and a resolution by objects of $\mathcal{I}$.

Remark 4.3. If $\mathfrak{a}$ is a dualizing k-variety, then both $\operatorname{mod}\mathfrak{a}$ and $\operatorname{mod}\mathfrak{a}^\circ$ are abelian.

Observation 4.4.

1. If Q is a strongly locally finite quiver (i.e. all the standard projectives are finitely generated and all the standard injectives are cofinitely generated) then the k-linear path category kQ of Q is a semi-hereditary dualizing k-variety.
2. Let $\mathfrak{a}$ be a dualizing k-variety, then $\operatorname{mod}\mathfrak{a}$ is also a dualizing k-variety ([2, Proposition 2.6]).
3. Let $\mathfrak{a}$ be a triangulated finite k-variety with Serre duality, then $\mathfrak{a}$ is a dualizing k-variety. The pseudokernels and pseudocokernels are given by the cones, and Serre duality shows that the category of standard injectives and the category of standard projectives are equivalent. (Moreover, a triangulated finite k-variety is a dualizing k-variety if and only if it satisfies Serre duality [12, Proposition 2.11])
4. It follows from [4] that any functorially finite full subcategory $\mathfrak{b}$ of a dualizing subcategory $\mathfrak{a}$ is again dualizing. This is the case when, for example, the full embedding $\mathfrak{b} \to \mathfrak{a}$ admits a left and a right adjoint (cf. Proposition 5.1).

Example 4.5. Let A_2 be the quiver $\cdot \to \cdot$, then the categories kA_2, $\operatorname{mod} kA_2$, and $D^b \operatorname{mod} kA_2$ are dualizing k-varieties.

Example 4.6. The category $\operatorname{coh}\mathbb{P}^1$ is not a dualizing k-variety since the standard projective module $\operatorname{Hom}(-, \mathcal{O}_{\mathbb{P}^1}) \in \operatorname{mod}(\operatorname{coh}\mathbb{P}^1)$ is not cofinitely generated. However, $D^b \operatorname{coh}\mathbb{P}^1$ has Serre duality and is thus a dualizing k-variety.

The next theorem is the main result of this section (compare with [10, Theorem 3.5] and [11, Theorem 1.4]).

Theorem 4.7. *Let $\mathfrak{a}$ be a finite k-variety. The following are equivalent:*

1. $\operatorname{mod}\mathfrak{a}$ *is abelian and has Serre duality,*
2. $\mathfrak{a}$ *is a dualizing k-variety and every object of* $\operatorname{mod}\mathfrak{a}$ *has finite projective and finite injective dimension.*

Proof. We denote by $\mathcal{P}$ and $\mathcal{I}$ the full additive subcategories of $\operatorname{Mod}\mathfrak{a}$ generated by the standard projectives and standard injectives, respectively.

First, assume $\mathfrak{a}$ is a dualizing k-variety and every object of $\operatorname{mod}\mathfrak{a}$ has finite projective and finite injective dimension. In this case, the canonical embeddings $K^b\mathcal{P} \to D^b \operatorname{Mod}_{\mathrm{lfd}}\mathfrak{a}$ and $K^b\mathcal{I} \to D^b \operatorname{Mod}_{\mathrm{lfd}}\mathfrak{a}$ induce equivalences of $K^b\mathcal{P}$ and $K^b\mathcal{I}$ with $D^b \operatorname{mod}\mathfrak{a}$. Let $N : K^b\mathcal{P} \to K^b\mathcal{I}$ be the equivalence induced by the Nakayama functor $\mathcal{P} \to \mathcal{I}$ given by sending $\mathfrak{a}(-, A)$ to $\mathfrak{a}(A, -)^*$. We define an autoequivalence $\mathbb{S} : D^b \operatorname{mod}\mathfrak{a} \to D^b \operatorname{mod}\mathfrak{a}$ by

$$\mathbb{S} : D^b \operatorname{mod}\mathfrak{a} \cong K^b\mathcal{P} \xrightarrow{N} K^b\mathcal{I} \cong D^b \operatorname{mod}\mathfrak{a}.$$

Every $X, Y \in D^b \operatorname{mod} \mathfrak{a}$ correspond to bounded complexes of projectives in $K^b\mathcal{P}$ and in order to show $\mathbb{S}$ is a Serre functor we may reduce to the case where $X \cong \mathfrak{a}(-,A)$ and $Y \cong \mathfrak{a}(-,B)$. The required isomorphism

$$\operatorname{Hom}(\mathfrak{a}(-,A), \mathfrak{a}(-,B)) \cong \operatorname{Hom}(\mathfrak{a}(-,B), \mathfrak{a}(A,-)^*)^*$$

follows from the Yoneda lemma.

For the other direction, we assume $\operatorname{mod} \mathfrak{a}$ is abelian and has Serre duality. For $A, B \in \operatorname{Ob} \operatorname{mod} \mathfrak{a}$, we have $\operatorname{Ext}^i_{\operatorname{mod} \mathfrak{a}}(A,B) \cong \operatorname{Hom}_{D^b \operatorname{mod} \mathfrak{a}}(A, B[i]) \cong \operatorname{Hom}_{D^b \operatorname{mod} \mathfrak{a}}(B[i], FA)^*$ such that $\operatorname{Ext}^i(A,-)$ can only be nonzero for finitely many i's. This shows that every object has finite projective dimension. Likewise one shows every object has finite injective dimension.

Since every object of $\operatorname{mod} \mathfrak{a}$ has a finite projective resolution, the natural embedding $K^b\mathcal{P} \to D^b \operatorname{mod} \mathfrak{a}$ is an equivalence. There is an equivalence $i : K^b\mathcal{I} \to D^b \operatorname{mod} \mathfrak{a}$ given by the composition

$$K^b\mathcal{I} \xrightarrow{N^{-1}} K^b\mathcal{P} \cong D^b \operatorname{mod} \mathfrak{a} \xrightarrow{\mathbb{S}} D^b \operatorname{mod} \mathfrak{a}$$

where $\mathbb{S}$ is a Serre functor on $D^b \operatorname{mod} \mathfrak{a}$. Since the isomorphisms

$$\operatorname{Hom}_{D^b \operatorname{mod} \mathfrak{a}}(\mathfrak{a}(-,B), \mathfrak{a}(A,-)^*) \cong \operatorname{Hom}_{D^b \operatorname{mod} \mathfrak{a}}(\mathfrak{a}(-,B), i\mathfrak{a}(A,-)^*)$$

are natural in $A, B \in \operatorname{Ob} \mathfrak{a}$, the Yoneda lemma implies that $i : K^b\mathcal{I} \to D^b \operatorname{mod} \mathfrak{a}$ is naturally equivalent to the canonical embedding. This shows that every object of $D^b \operatorname{mod} \mathfrak{a}$ has a finite resolution by standard injectives.

Since $\operatorname{mod} \mathfrak{a}$ is abelian, $\mathfrak{a}$ has pseudokernels. The canonical embedding $K^b\mathcal{I} \to D^b \operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$ induces an equivalence with $D^b \operatorname{mod} \mathfrak{a}$, and thus $\mathfrak{a}$ has pseudocokernels. Proposition 4.1 yields that $\mathfrak{a}$ is a dualizing k-variety. □

Corollary 4.8. *Let $\mathfrak{a}$ be a finite k-variety. The category* $\operatorname{mod} \mathfrak{a}$ *is abelian, hereditary, and has Serre duality if and only if $\mathfrak{a}$ is a semi-hereditary dualizing k-variety.*

Remark 4.9. The following example shows that there are dualizing k-varieties $\mathfrak{a}$ such that $\operatorname{mod} \mathfrak{a}$ has infinite global dimension but $\operatorname{mod} \mathfrak{a}$ still has Serre duality.

Example 4.10. Let Q be an A_∞ quiver with zig-zag orientation where every zig has one more arrow than the preceding zag, labeled as in Figure 2. We define a relation on Q by requiring the composition of any two arrows to be zero. The associated additive category $\mathfrak{b}$ is a dualizing k-variety by Proposition 4.1. Since every object of $\operatorname{mod} \mathfrak{b}$ has finite projective dimension, Theorem 4.7 yields that $\operatorname{mod} \mathfrak{b}$ has Serre duality. However, denoting by $S(v)$ the simple representation associated with the vertex $v \in Q$, we see that the projective dimension of $S(a^i_i)$ is i so that the global dimension of $\operatorname{mod} \mathfrak{b}$ is infinite.



FIGURE 2.

Remark 4.11. The category $\operatorname{mod} \mathfrak{b}$ is piecewise hereditary. Indeed, let $\mathfrak{a} = kQ$ (thus disregarding the conditions on the arrows in Example 4.10) then one may define a functor by

$$\begin{array}{rcl} i : \mathfrak{a} & \to & D^b \operatorname{mod} \mathfrak{b} \\ a^i_j & \mapsto & S(a^i_j)[-j] \\ b^i_j & \mapsto & S(b^i_j)[-j] \end{array}$$

The essential image of i forms a partial tilting set ([17]) in $D^b \operatorname{mod} \mathfrak{b}$ and hence lifts to a fully faithful and exact functor $i : D^b \operatorname{mod} \mathfrak{a} \to D^b \operatorname{mod} \mathfrak{b}$. Since $\mathfrak{a}$ generates $D^b \operatorname{mod} \mathfrak{b}$ as a triangulated category, i will be an equivalence of triangulated categories.

Example 4.12. Let Q be an A_∞^∞ quiver with linear orientation. We put a relation on Q by requiring that the composition of any two arrows is zero. The associated category kQ is a clearly a dualizing k-variety (it is equivalent to the category $D^b \operatorname{mod} kA_2$ from Example 4.5).

Since all indecomposables of $\operatorname{mod} kQ$ are either projective-injective or isomorphic to a standard simple, it is easy to check that $\operatorname{mod} kQ$ has left and right almost split sequences, but Theorem 4.7 shows that $D^b \operatorname{mod} kQ$ has no Serre functor since every standard simple representation has infinite projective dimension.

Remark 4.13. As in Remark 4.11, one shows that the category $\operatorname{mod} kQ$ from the previous example is piecewise hereditary. Here $\operatorname{mod} kQ$ is derived equivalent to $\operatorname{mod} kA_\infty^\infty$ where A_∞^∞ has linear orientation.

Let $\mathfrak{a}$ be a finite k-variety, and A an indecomposable object in $\mathfrak{a}$. A map $f : A \to M$ is called *right almost split* if every nonsplit map $A \to B$ factors through f. Dually, a map $g : N \to A$ is called *left almost split* if every nonsplit map $B \to A$ factors through g.

We will say a finite k-variety $\mathfrak{a}$ is *locally finite and locally discrete* if every indecomposable object A of $\mathfrak{a}$ admits a right almost split map $A \to M$ and a left almost split map $N \to A$.

Example 4.14. Let Q be a finite quiver. The path category kQ is locally discrete and locally finite.

Example 4.15. Let $\mathcal{P}$ be the poset $\mathbb{N} \cdot \{+\infty\}$. We may draw the Auslander-Reiten quiver of $k\mathcal{P}$ as

$$\begin{tikzcd} 0 \arrow[r] & 1 \arrow[r] & 2 \arrow[r] & 3 \arrow[r, phantom, "\cdots\cdots"] & +\infty \end{tikzcd}$$

It is clear that $k\mathcal{P}$ is not locally discrete since there is no left almost split map $N \to (+\infty)$. Not that $+\infty$ is an accumulation point of $\mathcal{P}$.

We now give an equivalent formulation of these properties.

Proposition 4.16. *A finite k-variety is locally finite and locally discrete if and only if all standard simples of $\mathfrak{a}$ are finitely presented and cofinitely presented. Furthermore, every dualizing k-variety is locally finite and locally discrete.*

Proof. Assume $\mathfrak{a}$ is locally finite and locally discrete. For an indecomposable $A \in \operatorname{ind} \mathfrak{a}$, let $N \to A$ be a left almost split map which gives rise to a map in $\operatorname{mod} \mathfrak{a}$

$$\mathfrak{a}(-, N) \longrightarrow \mathfrak{a}(-, A).$$

of which the cokernel is the standard simple S_A. Dually, one shows all standard simples are cofinitely presented.

Next, assume all standard simples are finitely and cofinitely presented. We prove that every indecomposable $A \in \operatorname{ind} \mathfrak{a}$ admits a left almost split map $N \to A$, for a certain object $N \in \operatorname{Ob} \mathfrak{a}$. Consider a presentation of S_A

$$Q \longrightarrow \mathfrak{a}(-, A) \longrightarrow S_A \longrightarrow 0.$$

We may write the projective Q as $\mathfrak{a}(-, N)$ and, since $S_A \cong \mathfrak{a}(-, A)/\operatorname{rad} \mathfrak{a}(-, A)$, the induced map $N \to A$ is left almost split. Dually, one proves a A admits a right almost split map $A \to M$.

For the last part, let $\mathfrak{a}$ be a dualizing k-variety and let $A \in \operatorname{Ob} \mathfrak{a}$ be an indecomposable object. Denote by $S_A \in \operatorname{Ob} \operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$ the associated standard simple representation. We know there is an epimorphism $\mathfrak{a}(-, A) \to S_A$ in $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$ and, since DS_A is a simple object in $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}$, there is an epimorphism $\mathfrak{a}(A, -) \to DS_A$ in $\operatorname{Mod}_{\mathrm{lfd}} \mathfrak{a}^\circ$. Applying D yields that S_A is the image of a map $f : \mathfrak{a}(-, A) \to \mathfrak{a}(A, -)^*$. Proposition 4.1 yields that $\ker f \in \operatorname{mod} \mathfrak{a}$.

There is now a short exact sequence $0 \to K \to \mathfrak{a}(-, A) \to S_A \to 0$ which shows $S_A \in \operatorname{Ob} \operatorname{mod} \mathfrak{a}$. □

Remark 4.17. That all standard simples representations of a dualizing k-variety are finitely presented, has already be shown in [2, Proposition 3.2].

5. (CO)REFLECTIVE SUBVARIETIES

Recall that a full replete (=closed under isomorphisms) subcategory of a category is called reflective or coreflective if the embedding has a left or a right adjoint, respectively.

In this section we will consider such reflective and coreflective subcategories of finite k-varieties. These subcategories occur often in the semi-hereditary case (Proposition 5.2) and the following proposition show they preserve the properties we are interested in.

Proposition 5.1. *Let $\mathfrak{a}$ be a dualizing k-variety, $\mathfrak{b}$ a finite k-variety, and $i : \mathfrak{b} \to \mathfrak{a}$ a full embedding admitting a left adjoint i_L and a right adjoint i_R. Then $\mathfrak{b}$ is also a dualizing k-variety. Furthermore, if (the abelian category) $\operatorname{mod} \mathfrak{a}$ satisfies Serre duality, then so does $\operatorname{mod} \mathfrak{b}$.*

Proof. Let $B_1 \to B_2$ be a map in $\mathfrak{b}$. Let $K, C \in \operatorname{Ob} \mathfrak{a}$ be a pseudokernel and pseudocokernel of the map $iB_1 \to iB_2$ in $\mathfrak{a}$, respectively. It is easily checked that $i_R K$ is a pseudokernel and $i_L C$ is a pseudocokernel of $B_1 \to B_2$.

Let $B \in \operatorname{Ob} \mathfrak{b}$ and let $\mathfrak{a}(-, A_1) \to \mathfrak{a}(-, A_0) \to \mathfrak{a}(iB, -)^* \to 0$ be a projective resolution in $\operatorname{mod} \mathfrak{a}$, then $\mathfrak{b}(-, i_R A_1) \to \mathfrak{b}(-, i_R A_0) \to \mathfrak{b}(B, -)^* \to 0$ is a projective resolution in $\operatorname{mod} \mathfrak{b}$. This shows that every standard injective in $\operatorname{mod} \mathfrak{b}$ is finitely presented. Likewise, one shows that every (standard) projective in $\mathfrak{b}$ is cofinitely presented. Proposition 4.1 yields that $\mathfrak{b}$ is a dualizing k-variety.

Assume now furthermore that $\operatorname{mod} \mathfrak{a}$ satisfies Serre duality. By Theorem 4.7, it suffices to show that every object of $\operatorname{mod} \mathfrak{b}$ has finite projective and finite injective dimension. Thus consider an $M \in \operatorname{mod} \mathfrak{b}$ and assume a projective resolution of $\mathfrak{a} \otimes_{\mathfrak{b}} M$ is given by

$$0 \to \mathfrak{a}(-, A_n) \to \cdots \to \mathfrak{a}(-, A_1) \to \mathfrak{a}(-, A_0) \to \mathfrak{a} \otimes_{\mathfrak{b}} M \to 0.$$

Restricting this to $\mathfrak{b}$ and using that $(\mathfrak{a} \otimes_{\mathfrak{b}} -)_{\mathfrak{b}}$ is equivalent to the identity functor on $\operatorname{mod} \mathfrak{b}$, we find

$$0 \to \mathfrak{b}(-, i_R A_n) \to \cdots \to \mathfrak{b}(-, i_R A_1) \to \mathfrak{b}(-, i_R A_0) \to M \to 0$$

such that M has finite projective dimension. Analogously, one shows that every object in $\operatorname{mod} \mathfrak{b}$ has a finite resolution by standard injectives. □

The following proposition gives some examples of reflective and/or coreflective subvarieties.

Proposition 5.2. *Let $\mathfrak{a}$ be a semi-hereditary finite k-variety.*

1. *Let $\mathcal{Z}$ be a set of objects of $\operatorname{mod} \mathfrak{a}$ such that $\sum_{Z \in \mathcal{Z}} \dim Z(A) < \infty$ for all $A \in \operatorname{Ob} \mathfrak{a}$ and $\operatorname{Ext}^1(Z_1, Z_2) = 0$ for all $Z_1, Z_2 \in \mathcal{Z}$. Define a full subcategory $\mathfrak{a}_{\perp \mathcal{Z}}$ of $\mathfrak{a}$ given by*
$$B \in \mathfrak{a}_{\perp \mathcal{Z}} \iff \forall Z \in \mathcal{Z} : Z(B) = 0.$$
The embedding $i : \mathfrak{a}_{\perp \mathcal{Z}} \to \mathfrak{a}$ has a left and right adjoint.
2. *Let $Y \in \operatorname{ind} \mathfrak{a}$. The embedding $\operatorname{supp}(-, Y) \to \mathfrak{a}$ has a left adjoint.*
3. *Let $X, Y \in \operatorname{ind} \mathfrak{a}$. If $\mathfrak{a}$ is a dualizing k-variety, then the embedding $[X, Y] \to \mathfrak{a}$ has a left and a right adjoint.*

Proof. (1) We start by defining the left adjoint $i_L : \mathfrak{a} \to \mathfrak{b}$. Let $A_1 \to A_2$ be a map in $\mathfrak{a}$ and consider the following commutative diagram with exact rows

$$\begin{tikzcd}
0 \arrow[r] & \mathfrak{a}(-, A_1') \arrow[r] \arrow[d, dashed] & \mathfrak{a}(-, A_1) \arrow[r] \arrow[d] & \bigoplus_{Z \in \mathcal{Z}} \operatorname{Hom}(\mathfrak{a}(-, A_1), Z)^* \otimes_k Z \arrow[d] \\
0 \arrow[r] & \mathfrak{a}(-, A_2') \arrow[r] & \mathfrak{a}(-, A_2) \arrow[r] & \bigoplus_{Z \in \mathcal{Z}} \operatorname{Hom}(\mathfrak{a}(-, A_2), Z)^* \otimes_k Z
\end{tikzcd}$$

Note that both $\bigoplus_{Z \in \mathcal{Z}} \operatorname{Hom}(\mathfrak{a}(-, A_1), Z)^* \otimes_k Z$ and $\bigoplus_{Z \in \mathcal{Z}} \operatorname{Hom}(\mathfrak{a}(-, A_2), Z)^* \otimes_k Z$ lie in $\operatorname{mod} \mathfrak{a}$ (since $\sum_{Z \in \mathcal{Z}} \dim Z(A) < \infty$ for all $A \in \operatorname{Ob} \mathfrak{a}$) and that $Z(A_1') = Z(A_2') = 0$ for all $Z \in \mathcal{Z}$ (since $\operatorname{Ext}^1(Z_1, Z_2) = 0$ for all $Z_1, Z_2 \in \mathcal{Z}$) so that $A_1', A_2' \in \operatorname{Ob} \mathfrak{b}$.

Since the kernel, the Yoneda embedding, and the tensor product are functorial, it is easy to see that the correspondence from $A_1 \to A_2$ to $A_1' \to A_2'$ is functorial. It is readily checked that this functor is right adjoint to the embedding.

The left adjoint is defined in a dual way.

(2) For every object $A \in \operatorname{Ob} \mathfrak{a}$, consider the canonical map $f_A : \mathfrak{a}(-,A) \to \mathfrak{a}(-,Y) \otimes \operatorname{Hom}(A,Y)^*$. Since $\operatorname{mod} \mathfrak{a}$ is hereditary, the image $\operatorname{im} f_A$ is a representable functor. Choose an object $A' \in \operatorname{Ob} \mathfrak{a}$ such that $\operatorname{im} f_A \cong \mathfrak{a}(-,A')$.

Let $A \to B$ be a map in $\mathfrak{a}$. The following commutative diagram

$$\begin{tikzcd}
\mathfrak{a}(-,A) \arrow[rr, "f"] \arrow[ddd] \arrow[dr] & & \mathfrak{a}(-,Y) \otimes \operatorname{Hom}(A,Y)^* \arrow[ddd] \\
& \mathfrak{a}(-,A') \arrow[ur] \arrow[d, dashed] & \\
& \mathfrak{a}(-,B') \arrow[dr] & \\
\mathfrak{a}(-,B) \arrow[rr, "g"] \arrow[ur] & & \mathfrak{a}(-,Y) \otimes \operatorname{Hom}(B,Y)^*
\end{tikzcd}$$

gives a map $\mathfrak{a}(-,A') \to \mathfrak{a}(-,B')$. The objects A' and B' lie in $\operatorname{supp} \mathfrak{a}(-,Y)$ and the correspondence $A \to B$ to $A' \to B'$ is functorial. It is readily checked that this defines a left adjoint $i_L : \mathfrak{a} \to \operatorname{supp} \mathfrak{a}(-,Y)$ to the functor i.

(3) We will only prove the embedding has a left adjoint; the right adjoint is dual. Let $\mathfrak{a}(-,Z)$ be a kernel of the canonical map $\mathfrak{a}(-,Y) \to \mathfrak{a}(X,-)^* \otimes \mathfrak{a}(X,Y)$. We show that for any $A \in \operatorname{supp} \mathfrak{a}(-,Y)$ we have $A \in \operatorname{supp} \mathfrak{a}(X,-)^*$ if and only if $\mathfrak{a}(A,Z) = 0$. Note that by construction of Z we already have that $\mathfrak{a}(X,Z) = 0$, so that $\mathfrak{a}(A,Z) \neq 0$ implies that $A \notin \operatorname{supp} \mathfrak{a}(X,-)^*$. For the other direction: if $\mathfrak{a}(A,Z) = 0$ then the exact sequence $0 \to \mathfrak{a}(-,Z) \to \mathfrak{a}(-,Y) \to \mathfrak{a}(X,-)^* \otimes \mathfrak{a}(X,Y)$ shows that $A \in \operatorname{supp} \mathfrak{a}(X,-)^*$.

By applying the first part of this proposition, we know that $[X,Y] \to \operatorname{supp} \mathfrak{a}(-,Y)$ has a left adjoint. The second part of the proposition implies $\operatorname{supp} \mathfrak{a}(-,Y) \to \mathfrak{a}$ has a left adjoint. This proves the final statement.

□

The following example shows that the conditions in the first statement of Proposition 5.2 are necessary.

Example 5.3. Let $\mathcal{L} = \mathbb{N} \cdot -\mathbb{N}$. By choosing $\mathcal{Z} = \{Z_i\}_{i \in \mathbb{N}}$ where

(1) $Z_i = k\mathcal{L}(-,i)$, or
(2) $Z_i = S_i$, where S_i is the simple representation associated with $i \in \mathbb{N}$

we find an associated subcategory $\mathfrak{b} = k\mathcal{L}_{\perp \mathcal{Z}}$ such that the natural embedding $\mathfrak{b} \to k\mathcal{L}$ does not have a left adjoint. Here the subcategory $\mathfrak{b}$ is generated by $-\mathbb{N}$.

Propositions 5.1 and 5.2 will be used in the next section to deduce properties of semi-hereditary dualizing k-varieties. We will give two examples to illustrate their use; a rigorous treatment follows in Proposition 6.2.

Example 5.4. Let $\mathfrak{a}$ be the semi-hereditary finite k-variety whose Auslander-Reiten quiver is given by



By removing $]X,Y[$ as in the first part of Proposition 5.2, we get a semi-hereditary finite k-variety with Auslander-Reiten quiver



which is clearly not locally discrete. We conclude that $\mathfrak{a}$ was not a *dualizing* k-variety.

Example 5.5. Let $\mathfrak{a}$ be the semi-hereditary finite k-variety whose Auslander-Reiten quiver is given by

$$\begin{tikzcd}
X \arrow[r] \arrow[d] & {} \arrow[r] \arrow[d] & {} \arrow[r] \arrow[d] & {} \arrow[r, dotted, no head] & {} \arrow[r] & Y \\
{} & {} & {} & \cdots & &
\end{tikzcd}$$

Again we remove $]X,Y[$ as in the first part of Proposition 5.2 and obtain a semi-hereditary finite k-variety with Auslander-Reiten quiver

$$\begin{tikzcd}
X \arrow[rrrr] \arrow[d] \arrow[rd] \arrow[rrd] \arrow[rrrd] & & & & Y \\
{} & {} & {} & {} & \cdots
\end{tikzcd}$$

which is not locally finite. Indeed the simple representation S_X is not cofinitely presented. Again we conclude $\mathfrak{a}$ is not a dualizing k-variety.

6. Threads

Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety. Recall from Proposition 4.16 that every indecomposable object $A \in \operatorname{Ob}\mathfrak{a}$ admits a right almost split map $A \to M$ and a left almost split map $N \to A$. Examples 5.4 and 5.5 show that most indecomposables lying in a convex subcategory should be –in some sense– nicely behaved. To make this rigorous, we introduce threads and thread objects.

Definition 6.1. If $\mathfrak{a}$ is a semi-hereditary dualizing k-variety, an indecomposable object $X \in \operatorname{ind}\mathfrak{a}$ will be called a *thread object* if X has a unique direct predecessor and a unique direct successor, or equivalently, there is a right almost split map $X \to M$ and a left almost split map $N \to X$ where M and N are indecomposable. In this case, we will denote the representatives of M and N in $\operatorname{ind}\mathfrak{a}$ by X^+ and X^-, respectively.

For $X, Y \in \operatorname{ind}\mathfrak{a}$, the subcategory $[X,Y]$ will be called a *thread* if every indecomposable object in $[X,Y]$ is a thread object in $\mathfrak{a}$. A thread is called *maximal* if it is not a proper subset of another thread. It is called *infinite* if it contains infinitely many nonisomorphic indecomposables.

An indecomposable object which is not a thread object is called a *nonthread object.*

Proposition 6.2. *Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety, and $X, Y, Z \in \operatorname{ind}\mathfrak{a}$, then*

1. $\operatorname{supp}\mathfrak{a}(X,-)^*$ *and* $\operatorname{supp}\mathfrak{a}(-,X)$ *have only finitely many nonthread objects.*
2. *if $[X,Y]$ is a thread, then* $\dim\mathfrak{a}(X,Y) = 1$*, and*
3. *if $[X,Y]$ and $[X,Z]$ are threads, then either $[X,Y] \subseteq [X,Z]$ or $[X,Z] \subseteq [X,Y]$.*

Proof. (1) It follows from Proposition 4.1 that $\mathfrak{a}(-,X)$ is cofinitely presented. Since $\mathfrak{a}$ is semi-hereditary, we see that $\operatorname{supp}\mathfrak{a}(-,X)$ has only finitely many nonthread objects. The other case is dual.

(2) Write $V = \mathfrak{a}(X,Y)^*$ and consider the canonical map $\mathfrak{a}(-,Y) \otimes_k V \to \mathfrak{a}(X,-)^*$ with image $F \in \operatorname{mod}\mathfrak{a}$. We have the following short exact sequences

$$0 \longrightarrow \mathfrak{a}(-,A) \longrightarrow \mathfrak{a}(-,Y) \otimes V \longrightarrow F \longrightarrow 0 \tag{2}$$

$$0 \longrightarrow F \longrightarrow \mathfrak{a}(X,-)^* \longrightarrow \mathfrak{a}(B,-)^* \longrightarrow 0 \tag{3}$$

It is clear that every direct summand of A maps nonzero to Y. Furthermore, from the second short exact sequence we obtain $\dim F(X) = 1$ so that $\dim V > 1$ would imply $\dim\mathfrak{a}(X,A) \neq 0$. Thus at least one indecomposable direct summand A_1 of A lies in $[X,Y[$, hence A_1 is a thread object with direct successor $A_1^+ \in \operatorname{ind}]X,Y] \subset \operatorname{ind}\mathfrak{a}$. It follows from the first short exact sequence that $\dim F(A_1) < \dim F(A_1^+)$.

However, applying $\operatorname{Hom}(\mathfrak{a}(-,Y),-)$ to the sequence (3) yields $\mathfrak{a}(B,Y) = 0$. In particular, since $\mathfrak{a}$ is semi-hereditary, we have for every $Z \in [X,Y]$ that $\dim\mathfrak{a}(B,Z) = 0$ and thus $\dim F(Z) = \dim\mathfrak{a}(X,Z)$. We find $\dim\mathfrak{a}(X,A_1) < \dim\mathfrak{a}(X,A_1^+)$, a contradiction since $\mathfrak{a}$ is semi-hereditary.

(3) Let $i : [X, Y] \to \mathfrak{a}$ be the natural embedding. By Proposition 5.2 we know that the embedding has a right adjoint $i_R : \mathfrak{a} \to [X, Y]$. Let A be an indecomposable direct summand of $i_R(Z)$.

If $A \cong Y$, then $\mathfrak{a}(Y, Z) \neq 0$ so that $[X, Y] \subseteq [X, Z]$.

Therefore, we will assume that $A \ncong Y$ and show that $iA \cong Z$. In this case, we have $A^+ \in \operatorname{ind}[X, Y]$. Since $A \to A^+$ is left and right almost split, we know that $\dim \mathfrak{a}(iA, Z) = \dim \mathfrak{a}(iA^+, Z)$ if $Z \ncong iA$. Using adjointness, we find $\dim \operatorname{Hom}(A, i_R Z) = \dim \operatorname{Hom}(A^+, i_R Z)$. This last statement is impossible since A is a direct summand of $i_R Z$. We find that $iA \cong Z$ and thus $\mathfrak{a}(Z, Y) \neq 0$. We conclude that $[X, Z] \subseteq [X, Y]$

□

Corollary 6.3. *Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety. Every thread $[X, Y]$ is equivalent to $k\mathcal{L}$ for a linearly ordered locally discrete poset $\mathcal{L}$ with a maximal and a minimal element.*

Proof. Write $\mathfrak{b} = [X, Y]$ and let $\mathcal{L} = \operatorname{ind}[X, Y]$. We define a poset structure on $\mathcal{L}$ by

$$A \leq B \iff \mathfrak{a}(A, B) \neq 0.$$

It follows from Proposition 6.2 that $\mathcal{L}, \leq$ is a linearly ordered locally discrete poset with unique minimal and maximal element. We will define a functor $k\mathcal{L} \to \mathfrak{b}$ as follows. For every $A \in \mathfrak{b}$, choose a nonzero element $b_{X,A} \in \mathfrak{a}(X, A)$.

For every $A, B \in \operatorname{ind} \mathfrak{b}$ with $\mathfrak{b}(A, B) \neq 0$, this gives a unique element $b_{A,B} \in \mathfrak{b}(A, B)$ with $b_{X,B} = b_{X,A} \circ b_{A,B}$.

The functor $k\mathcal{L} \to \mathfrak{b}$, which is the identity on objects and sending $(A, B) \in k\mathcal{L}(A, B)$ to $b_{A,B} \in \mathfrak{b}(A, B)$, is fully faithful and essentially surjective, and hence an equivalence. □

Corollary 6.4. *A connected semi-hereditary dualizing k-variety has only countably many sinks or sources.*

Corollary 6.5. *Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety, and let $[X, Y]$ be an thread. If $Z \in \operatorname{ind} \mathfrak{a}$ does not lie on $[X, Y]$, then every map $f : X \to Y$ factors through a nonzero map $X \to Y^+$.*

Proof. Let $\mathfrak{b}$ be the full subcategory of $\mathfrak{a}$ generated by objects not supported on $[X, Y]$. By Proposition 5.2, the embedding $i : \mathfrak{b} \to \mathfrak{a}$ has a left adjoint $i_L : \mathfrak{a} \to \mathfrak{b}$.

Using Proposition 6.2, it is straightforward to check that $i_L X \cong Y^+$ and $\dim \operatorname{Hom}(X, Y^+) = 1$.

For a $Z \in \mathfrak{b}$, every morphism in $\operatorname{Hom}(X, Z)$ factors through $\eta(1_X) \in \operatorname{Hom}(i \circ i_L X, Z)$ where $\eta : 1 \to i \circ i_L$ is the unit of the adjunction (i_L, i). □

Corollary 6.6. *Every thread in a semi-hereditary dualizing k-variety lies inside a maximum thread.*

Proof. Let $[X, Y]$ be a thread in a semi-hereditary dualizing k-variety $\mathfrak{a}$. Since $\mathfrak{a}(-, Y)$ is cofinitely presented, there is at least one sink S with $\mathfrak{a}(S, Y) \neq 0$ such that $\operatorname{supp} \mathfrak{a}(-, Y)$ has nonthread objects.

By Proposition 6.2 there are only finitely many nonthread objects in $\operatorname{supp} \mathfrak{a}(-, Y)$ and by Proposition 3.1 there are no cycles. We can thus find a nonthread object $Z \in \operatorname{supp} \mathfrak{a}(-, Y)$ such that Z is the only nonthread object in $[Z, Y]$. Write $Z^+ = X'$. It follows from Proposition 6.2 that $[X', Y]$ is a thread containing $[X, Y]$.

Dually, one finds a thread object object Y' such that $[X', Y']$ is a thread containing $[X', Y]$ and such that Y'^+ is a nonthread object. The thread $[X', Y']$ is the required maximal thread. □

Proposition 6.7. *Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety and let $\{[X_i, Y_i]\}_{i \in I}$ be a set of maximal threads. Let $\mathfrak{b}$ be the full subcategory of $\mathfrak{a}$ consisting of all objects without direct summands in one of these threads. Then $\mathfrak{b}$ is a semi-hereditary dualizing k-variety and the embedding $i : \mathfrak{b} \to \mathfrak{a}$ has both a left and a right adjoint.*

Proof. By Propositions 3.1 and 5.1 it suffices to show i has both a left and a right adjoint. Proposition 6.2 yields these threads are supported by disjoint subsets of $\operatorname{ind} \mathfrak{a}$ so that we may invoke Proposition 5.2, finishing the proof. □



FIGURE 3. An example of a thread quiver (left) and the underlying quiver (right)

7. CLASSIFICATION BY THREAD QUIVERS

Proposition 7.1. *Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety such that $[X,Y]$ is finite for all $X, Y \in \operatorname{ind} \mathfrak{a}$. Write Q for the Auslander-Reiten quiver of $\mathfrak{a}$, then $\mathfrak{a}$ is equivalent to kQ. Moreover, Q is a strongly locally finite quiver.*

Proof. For every two indecomposables $A, B \in \operatorname{ind} \mathfrak{a}$, we will choose $\dim_k \operatorname{Irr}(A,B)$ linearly independent irreducible maps $A \to B$. This gives a map $f : Q_0 \to \operatorname{Mor} \mathfrak{a}$, mapping the arrows in Q to the chosen irreducible maps in $\mathfrak{a}$.

Let $A, B \in \operatorname{ind} \mathfrak{a}$. Since every interval $[A,B]$ is assumed to be finite and no object in $\mathfrak{a}$ has nontrivial endomorphisms, every map $A \to B$ is a linear combination of compositions of chosen irreducible maps.

There is an obvious functor $F : kQ \to \mathfrak{a}$ which is fully faithful and essentially surjective. □

The previous proposition does not hold without the assumption on the intervals $[X,Y]$. However, in the case of dualizing k-varieties, these intervals are nicely behaved (Proposition 6.2) such that an analogue of Proposition 7.1 may be proven without any assumption on the intervals $[X,Y]$, by replacing the quiver Q by a thread quiver.

A *thread quiver* consists of the following information:

- A quiver $Q = (Q_0, Q_1)$ where Q_0 is the set of vertices and Q_1 is the set of arrows.
- A decomposition $Q_1 = Q_s \coprod Q_t$. Arrows in Q_s will be called *standard arrows*, while arrows in Q_t will be referred to as *thread arrows.*
- With every thread arrow α, there is an associated linearly ordered set $\mathcal{T}_\alpha$, possibly empty.

In drawing a thread quiver (cf. Figure 3), standard arrows will be represented by $\bullet \longrightarrow \bullet$, while thread arrows will be drawn as $\bullet \dashrightarrow \bullet$ labeled by the corresponding ordered set $\mathcal{T}$. If $\mathcal{T} = \{1,2,3,\ldots,n\}$ with the normal ordering, then we will only write n as a label; if $\mathcal{T} = \emptyset$, then no label will be written.

Every thread quiver Q has an underlying quiver where there is no distinction between the standard and the thread arrows, and where all arrows are unlabeled. To avoid confusion, we will refer to this underlying quiver by Q_r.

Starting from a semi-hereditary dualizing k-variety, we will associate a thread quiver in the following way.

Construction 7.2. Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety. Let $\{[X_i, Y_i]\}_{i \in I}$ be the set of all maximal *infinite* threads in $\mathfrak{a}$. The full subcategory of $\mathfrak{a}$ consisting of all objects which are not supported on $\bigcup_{i \in I} \operatorname{ind}[X_i, Y_i]$ is called $\mathfrak{b}$. By Proposition 6.7, this is again a semi-hereditary dualizing k-variety. Proposition 7.1 yields $\mathfrak{b} \cong kQ_r$ for a strongly locally finite quiver Q_r.

According to Corollary 6.3, every thread $[X_i, Y_i] \subset \operatorname{ind} \mathfrak{a}$ corresponds to a poset of the form $\mathbb{N} \cdot (\mathcal{T} \overset{\rightarrow}{\times} \mathbb{Z}) \cdot -\mathbb{N}$ where $\mathcal{T}$ is a linearly ordered set. We will replace one arrow from $X^- \longrightarrow Y^+$ in Q_r by $X^- \overset{\mathcal{T}}{\dashrightarrow} Y^+$.

Example 7.3. Let $\mathcal{L} = \mathbb{N} \cdot (\mathcal{T} \overset{\rightarrow}{\times} \mathbb{Z}) \cdot -\mathbb{N}$, then the thread quiver associated with $k\mathcal{L}$ is

$$\cdot \overset{\mathcal{T}}{\dashrightarrow} \cdot$$

Given a thread quiver Q, we will construct an associated semi-hereditary finite k-variety.

Construction 7.4. Let Q be a thread quiver with underlying strongly locally finite quiver Q_r. With every thread $t \in Q_t$, we denote by $f^t : k(\cdot \to \cdot) \longrightarrow kQ_r$ the functor associated with the obvious embedding $(\cdot \to \cdot) \longrightarrow Q_r$. We define the functor

$$f : \bigoplus_{t \in Q_t} k(\cdot \to \cdot) \longrightarrow kQ_r.$$

With every thread t, there is an associated linearly ordered set $\mathcal{T}_t$. We will write $\mathcal{L}_t = \mathbb{N} \cdot (\mathcal{T}_t \overset{\rightarrow}{\times} \mathbb{Z}) \cdot -\mathbb{N}$ and denote by

$$g^t : k(\cdot \to \cdot) \longrightarrow k\mathcal{L}_t$$

a chosen fully faithful functor given by mapping the extremal points of $\cdot \to \cdot$ to the minimal and maximal objects of $\mathcal{L}$, respectively. We will write

$$g : \bigoplus_{t \in Q_t} k(\cdot \to \cdot) \longrightarrow \bigoplus_{t \in Q_t} k\mathcal{L}_t.$$

We define the category kQ as a 2-pushout in $k\mathfrak{AdCat}$ of the following diagram.

$$\begin{tikzcd}
\bigoplus_{t \in Q_t} k(\cdot \to \cdot) \arrow[r, "f"] \arrow[d, "g"'] & kQ_r \arrow[d, dashed, "i"] \\
\bigoplus_{t \in Q_t} k\mathcal{L}_t \arrow[r, dashed, "j"'] & kQ
\end{tikzcd}$$

Remark 7.5. The above 2-pushout does indeed exist since 2-colimits exist in the 2-category $\mathfrak{Cat}$ and the functor $k : \mathfrak{Cat} \to k\mathfrak{AdCat}$ has a right adjoint (cf. §2.4).

Remark 7.6. Given a thread quiver Q, Construction 7.4 will define kQ only up to equivalence.

Example 7.7. Let Q be the thread quiver given by



The associated finite k-variety is equivalent to $k(\mathbb{N} \cdot \mathcal{T} \overset{\rightarrow}{\times} \mathbb{Z} \cdot -\mathbb{N})$.

Remark 7.8. Different thread quivers Q, Q' may give rise to equivalent categories kQ and kQ' as shown in the following example.

Example 7.9. The following three thread quivers give rise to equivalent categories.







Observation 7.10. The functors f^t have left adjoints (denoted by f^t_L) and right adjoints (denoted by f^t_R). The induced functors

$$f_L, f_R : kQ_r \longrightarrow \bigoplus_{t \in Q_t} k(\cdot \to \cdot)$$

are left and right adjoints for f, respectively.

Similarly, the functors g_t have left adjoints (denoted by g^t_L) and right adjoints (denoted by g^t_R, giving rise to the functors

$$g_L, g_R : \bigoplus_{t \in Q_t} k\mathcal{L}_t \longrightarrow \bigoplus_{t \in Q_t} k(\cdot \to \cdot)$$

which are then left and right adjoints for g, respectively.

The functor g is fully faithful and its adjoints g_L and g_R are faithful. Note that f^t will always be faithful, but in general not full. Its adjoints f^t_L and f^t_R are in general not full nor faithful.

Example 7.11. The functor f^t is not full when Q is given by the following thread quiver.



In what follows it will be convenient to assume the functors f^t are fully faithful. For this we will use the following lemma.

Lemma 7.12. *Let Q be a thread quiver. There is a thread quiver Q' such that*

1. *$kQ \cong kQ'$, and*
2. *the functors ${f'}^t : k(\cdot \to \cdot) \longrightarrow kQ'_r$ from Construction 7.4 are fully faithful.*

Proof. Define a thread quiver Q' by replacing all thread arrows $x \overset{\mathcal{T}}{\dashrightarrow} y$ in the thread quiver Q by $x \longrightarrow \cdot \overset{\mathcal{T}}{\dashrightarrow} \cdot \longrightarrow y$. Since the associated functors ${f'}^t : k(\cdot \to \cdot) \longrightarrow kQ'_r$ are fully faithful, we need only to prove $kQ \cong kQ'$.

We start by defining a 2-pushout

$$\begin{tikzcd}
\bigoplus_{t \in Q'_t} k(a_t \to b_t \to c_t \to d_t) \arrow[r, "f_1"] \arrow[d, "g_1"'] & kQ'_r \arrow[d, dashed] \\
\bigoplus_{t \in Q'_t} k\mathcal{L}_t \arrow[r, dashed] & \mathfrak{p}
\end{tikzcd}$$

in $k\mathfrak{AdCat}$ where the maps are given by the following descriptions. The functor ${f_1}^t$, for each thread $t : x_t \overset{\mathcal{T}}{\dashrightarrow} y_t$ in Q', is given as a faithful functor which maps b_t and c_t to x_t and y_t respectively, and a_t and d_t to the direct predecessor of x_t and the direct successor of y_t, respectively.

The functor g_1^t is a fully faithful functor given by

$$\begin{array}{rcl}
a_t & \mapsto & 0_t \\
b_t & \mapsto & 1_t \\
c_t & \mapsto & -1_t \\
d_t & \mapsto & -0_t
\end{array}$$

where 0_t and -0_t are the minimal and maximal element of $\mathcal{L}_t$ respectively, 1_t is the direct successor of 0_t, and -1_t is the direct predecessor of -0_t.

Define a functor

$$H_1 : \bigoplus_{t \in Q_t} k(x_t \to y_t) \longrightarrow \bigoplus_{t \in Q_t} k(a_t \to b_t \to c_t \to d_t)$$

by mapping x_t to a_t, y_t to d_t, and the generator $x_t \to y_t$ to the composition $a_t \to b_t \to c_t \to d_t$. We also consider the natural embedding of categories $F : kQ_r \longrightarrow kQ'_r$.

Consider the following diagram where we write A_2 and A_4 for $(\cdot \to \cdot)$ and $(\cdot \to \cdot \to \cdot \to \cdot)$ respectively.

$$\begin{tikzcd}
\bigoplus_t kA_2 \arrow[r, "f"] \arrow[d, "H_1"'] & kQ_r \arrow[d, "F"'] \arrow[dd, bend left=60, "i"] \\
\bigoplus_t kA_4 \arrow[r, "f_1"] \arrow[d, "g_1"'] & kQ'_r \\
\bigoplus_t k\mathcal{L}_t \arrow[r, "j"'] & kQ
\end{tikzcd}$$

where the outer diagram is a 2-pushout. It is readily verified that the upper square is also a 2-pushout. The 2-natural transformation given by $i : kQ_r \to kQ$ and $j : \bigoplus_{t \in Q_t} k\mathcal{L}_t \to kQ$ induces a functor $i' : kQ'_r \to kQ$. Using the universal property, it is now straightforward to check that the lower square is also a 2-pushout. This shows that $\mathfrak{p} \cong kQ$.

To show that $\mathfrak{p} \cong kQ'$, consider the 2-pushout

$$\begin{tikzcd}
\bigoplus_{t\in Q'_t} k(x'_t \to y'_t) \arrow[r, "f'"] \arrow[d, "g'"'] & kQ'_r \arrow[d, dashed, "i'"] \\
\bigoplus_{t\in Q'_t} k\mathcal{L}'_t \arrow[r, dashed, "j'"] & kQ'
\end{tikzcd}$$

where $\mathcal{L}'_t = \mathbb{N}_0 \cdot (\mathcal{T} \overset{\to}{\times} \mathbb{Z}) \cdot -\mathbb{N}_0$ and $\mathbb{N}_0 = \mathbb{N} \setminus \{0\}$. Note that $\mathcal{L}'_t \cong \mathcal{L}_t$.

We define a functor

$$H_2 : \bigoplus_{t\in Q'_t} k(x'_t \to y'_t) \longrightarrow \bigoplus_{t\in Q'_t} k(a_t \to b_t \to c_t \to d_t)$$

by mapping x'_t to b_t, y'_t to c_t, and the generator $x_t \to y_t$ to the generator $b_t \to c_t$. We also consider the natural embedding of categories $G : \bigoplus_{t\in Q'_t} k\mathcal{L}'_t \longrightarrow \bigoplus_{t\in Q'_t} k\mathcal{L}_t$.

As before, one obtains a diagram

$$\begin{tikzcd}
\bigoplus_t kA_2 \arrow[r, "H_2"] \arrow[d, "g'"'] & \bigoplus_t kA_4 \arrow[r, "f_1"] \arrow[d, "g_1"'] & kQ'_r \arrow[d, "i"] \\
\bigoplus_t k\mathcal{L}'_t \arrow[r, "G"] \arrow[rr, bend right, "j'"'] & \bigoplus_t k\mathcal{L}_t & kQ'
\end{tikzcd}$$

where the left square and the outer diagram are 2-pushouts. Using the universal property, one shows that the right square is also a 2-pushout. This shows that $\mathfrak{p} \cong kQ'$ and thus $kQ \cong kQ'$. □

Example 7.13. Let Q be the quiver of Example 7.11. The quiver Q' constructed in the proof of Lemma 7.12 is given by



Remark 7.14. Let thread quiver Q' will be strongly locally finite if and only if Q is.

Proposition 7.15. *The functor i from Construction 7.4 is fully faithful and has both a left and a right adjoint. If f^s is fully faithful for a thread $s \in Q_t$, then the functor j^s is fully faithful and has both a left and a right adjoint.*

Proof. Since g is fully faithful and has a left and a right adjoint, Theorem A.1 shows the same properties hold for i.

Let $s \in Q_t$ be a thread such that $f^s : k(\cdot \to \cdot) \longrightarrow kQ_r$ is fully faithful. We will construct the left adjoint of $j^s : k\mathcal{L}_s \to kQ$. First, consider the 2-pushouts

$$\begin{tikzcd}
k(\cdot \to \cdot) \arrow[r, "f^s"] \arrow[d, "g^s"'] & kQ_r \arrow[d, dashed, "G"] \arrow[dl, phantom, "\overset{\alpha}{\Nearrow}"] \\
k\mathcal{L}_s \arrow[r, dashed, "F^s"'] & \mathfrak{p}
\end{tikzcd}
\qquad
\begin{tikzcd}
\bigoplus_{t\neq s} k(\cdot \to \cdot) \arrow[r, "(G\circ f^t)_t"] \arrow[d, "(g^t)_t"'] & \mathfrak{p} \arrow[d, dashed, "i'"] \arrow[dl, phantom, "\overset{\beta}{\Nearrow}"] \\
\bigoplus_{t\neq s} k\mathcal{L}_t \arrow[r, dashed] & \mathfrak{p}'
\end{tikzcd}$$

Using the universal property, it is straightforward to proof $\mathfrak{p}' \cong kQ$ and that $j^s = i' \circ F^s$. Since both f^s and $(g^t)_{t\neq s}$ are fully faithful and have a left and a right adjoint, Theorem A.1 yields that the same properties hold for F^s and i', and hence also for j^s. □

Although Construction 7.4 will not yield all (semi-hereditary) finite k-varieties, the following theorem says it is general enough to give all semi-hereditary dualizing k-varieties.

Proposition 7.16. *If Q is a thread quiver with Q_r strongly locally finite, then kQ is a semi-hereditary dualizing k-variety.*

Proof. We may assume the quiver Q satisfies the property of Lemma 7.12 such that Proposition 7.15 yields i and j^t have right adjoints, for all threads $t \in Q_t$. In this case, a finitely generated projective module $kQ(-,A) \in \operatorname{Mod} kQ$ corresponds to a diagram

$$\begin{tikzcd}
\bigoplus_t k(\cdot \to \cdot) \arrow[r, "f"] \arrow[d, "g"'] \arrow[dr, phantom, "{}^{\alpha}\Nearrow"] & kQ_r \arrow[d, "i"] \arrow[ddr, bend left, "kQ_r(-{,}i_RA)"] & \\
k\mathcal{L} \arrow[r, "j"'] \arrow[drr, bend right, "\mathcal{L}(-{,}j_RA)"'] \arrow[dr, phantom, "{}^{\beta}\Uparrow"] & kQ \arrow[dr, dashed, "kQ(-{,}A)"] \arrow[r, phantom, "\Swarrow_{\gamma}"] & {} \\
& & \operatorname{Mod} k
\end{tikzcd}$$

where we have written $k\mathcal{L} = \bigoplus_t k\mathcal{L}_t$. This shows that the natural embedding $\operatorname{Mod} kQ \to \operatorname{Mod} kQ_r \oplus \operatorname{Mod} k\mathcal{L}$ restricts to an embedding $\operatorname{mod} kQ \to \operatorname{mod} kQ_r \oplus \operatorname{mod} k\mathcal{L}$. Since the latter category is Hom-finite, so is the former. In particular we know that kQ is Hom-finite. It follows easily from Lemma 7.17 below that idempotents split in kQ, such that kQ is a finite k-variety.

To show that kQ is semi-hereditary (thus that $\operatorname{mod} kQ$ is abelian and hereditary) it suffices to show that the kernel between finitely generated projectives of $\operatorname{Mod} kQ$ is projective and splits off.

Let $kQ(-,A) \to kQ(-,B)$ be a map between finitely generated projectives. Since $\operatorname{mod} kQ_r$ and $\operatorname{mod} k\mathcal{L}$ are hereditary, we get the exact sequences

$$\begin{tikzcd}
0 \arrow[r] & kQ_r(-,K_1) \arrow[r] & kQ_r(-,i_RA) \arrow[r] & kQ_r(-,i_RB)
\end{tikzcd}$$

and

$$\begin{tikzcd}
0 \arrow[r] & k\mathcal{L}(-,K_2) \arrow[r] & k\mathcal{L}(-,j_RA) \arrow[r] & k\mathcal{L}(-,j_RB)
\end{tikzcd}$$

where the kernels split off.

These exact sequences induce the following commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & kQ_r(f-,K_1) \arrow[r] \arrow[d, dashed] & kQ_r(f-,i_RA) \arrow[r] \arrow[d] & kQ_r(f-,i_RB) \arrow[d] \\
0 \arrow[r] & k\mathcal{L}(g-,K_2) \arrow[r] & k\mathcal{L}(g-,j_RA) \arrow[r] & k\mathcal{L}(g-,j_RB)
\end{tikzcd}$$

which shows that the kernel of $kQ(-,A) \to kQ(-,B)$ splits off, and hence is projective. We have shown kQ is semi-hereditary.

We proceed to show kQ is a dualizing k-variety, thus that $D : \operatorname{Mod}_{\mathrm{lfd}} kQ \to \operatorname{Mod}_{\mathrm{lfd}} kQ^{\circ}$ induces a duality $D : \operatorname{mod} kQ \to \operatorname{mod} kQ^{\circ}$. Let $M \in \operatorname{mod} kQ$ giving the following diagram

$$\begin{tikzcd}
\bigoplus_t k(\cdot \to \cdot) \arrow[r, "f"] \arrow[d, "g"'] \arrow[dr, phantom, "{}^{\alpha}\Nearrow"] & kQ_r \arrow[d, "i"] \arrow[ddr, bend left, "N"] & \\
k\mathcal{L} \arrow[r, "j"'] \arrow[drr, bend right, "L"'] \arrow[dr, phantom, "{}^{\beta}\Uparrow"] & kQ \arrow[dr, dashed, "M"'] \arrow[r, phantom, "\Swarrow_{\gamma}"] & {} \\
& & \operatorname{mod} k
\end{tikzcd}$$

Composing with the functor $\operatorname{Hom}(-,k) : \operatorname{mod} k \to \operatorname{mod} k$, gives the diagram

$$\begin{tikzcd}
\bigoplus_t k(\cdot \to \cdot) \arrow[r, "f"] \arrow[d, "g"'] \arrow[dr, phantom, "\alpha\Uparrow"] & kQ_r \arrow[d, "i"] \arrow[ddr, bend left, "DN"] & \\
k\mathcal{L} \arrow[r, "j"'] \arrow[drr, bend right, "DL"'] & kQ \arrow[dr, dashed, "DM"'] \arrow[r, phantom, "\Swarrow"] \arrow[d, phantom, "\Uparrow"] & \\
 & & \operatorname{mod} k
\end{tikzcd}$$

Since kQ_r and $k\mathcal{L}$ are dualizing k-varieties, we see that $DM \in \operatorname{mod} kQ^\circ$. Likewise, one shows that the functor $D : \operatorname{Mod}_{\mathrm{lfd}} kQ^\circ \to \operatorname{Mod}_{\mathrm{lfd}} kQ$ restricts to $D : \operatorname{mod} kQ^\circ \to \operatorname{mod} kQ$, and hence induces a duality. $\square$

We have used the following lemma.

Lemma 7.17. *Let $\mathfrak{a} : \mathcal{I} \to k\mathfrak{AdCat}$ be a 2-functor. If for every $i \in \operatorname{Ob}\mathcal{I}$, all idempotents split in $\mathfrak{a}(i)$, then all idempotents split in $2\operatorname{colim}\mathfrak{a}$.*

Proof. Recall ([8]) that a category in which all idempotents split is called Cauchy complete. Denote by $\mathfrak{Cc}.\mathfrak{Cat}$ the 2-category of small Cauchy complete k-linear additive categories. It follows from [8, Proposition 6.5.9] that the embedding $\mathfrak{Cc}.\mathfrak{Cat} \to k\mathfrak{AdCat}$ has a left adjoint $k\mathfrak{AdCat} \to \mathfrak{Cc}.\mathfrak{Cat}$. Hence taking Cauchy completions commute with 2-colimits ([9, Proposition 6.7.3] or [13, §3.2]). $\square$

We are now ready to complete the classification of semi-hereditary dualizing k-varieties by thread quivers.

Theorem 7.18. *Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety, then $\mathfrak{a}$ may be obtained by Construction 7.4 from a thread quiver Q where Q is strongly locally finite.*

Proof. It has been established in Proposition 7.16 that kQ is indeed a semi-hereditary dualizing k-variety, so that we need only to show every semi-hereditary dualizing k-variety arises in this way (up to equivalence).

Let $\mathfrak{a}$ be a semi-hereditary dualizing k-variety and let Q be the corresponding thread quiver as in Construction 7.2. There are obvious embeddings $i^{\mathfrak{a}} : kQ_r \to kQ$ and $j^{t,\mathfrak{a}} : k\mathcal{L}_t \to \mathfrak{a}$ which give a functor $F : kQ \to \mathfrak{a}$ as in the following diagram

$$\begin{tikzcd}
\bigoplus_t k(\cdot \to \cdot) \arrow[r, "f"] \arrow[d, "g"'] \arrow[dr, phantom, "\alpha\Uparrow"] & kQ_r \arrow[d, "i"] \arrow[ddr, bend left, "i^{\mathfrak{a}}"] & \\
\oplus_t k\mathcal{L}_t \arrow[r, "j"'] \arrow[drr, bend right, "j^{\mathfrak{a}}"'] & kQ \arrow[dr, dashed, "F"'] \arrow[r, phantom, "\Swarrow"] \arrow[d, phantom, "\Uparrow"] & \\
 & & \mathfrak{a}
\end{tikzcd}$$

Since the induced functor $kQ_r \oplus k\mathcal{L} \to \mathfrak{a}$ is essentially surjective, so is the functor F. We will continue by showing F is also fully faithful by using the description of kQ given in §2.3.

Since $i^{\mathfrak{a}} \cong F \circ i$ and both i and $i^{\mathfrak{a}}$ are fully faithful, F induces a bijection $kQ(iX, iY) \to \mathfrak{a}(FiX, FiY)$.

Next, consider the embedding $G^t : k\mathcal{L}'_t \to k\mathcal{L}_t$ as in the proof of Lemma 7.12, thus the only indecomposables of $k\mathcal{L}_t$ which do not lie in $k\mathcal{L}'_t$ are the minimal and maximal elements of $\mathcal{L}_t$. The essential image of $j'^{t,\mathfrak{a}} = j^{t,\mathfrak{a}} \circ G^t$ is a maximal thread in $\mathfrak{a}$. Moreover, $j'^{t,\mathfrak{a}}$ is faithful and Proposition 6.2 shows it is full as well.

As above we find that F induces a bijection $kQ(j^tX, j^tY) \to \mathfrak{a}(Fj^tX, Fj^tY)$ for all X, Y in the essential image of $k\mathcal{L}'_t$.

Following Corollary 6.5 and its dual, this suffices to show that $F : kQ \to \mathfrak{a}$ is indeed a fully faithful functor. We have shown that F is an equivalence. $\square$

8. Representations of thread quivers

Let Q be a thread quiver. We define the categories $\operatorname{Rep}_k Q$ and $\operatorname{rep}_k Q$ to be $\operatorname{Mod} kQ$ and $\operatorname{mod} kQ$, respectively. We have the following result.

Theorem 8.1. *Let $\mathcal{A}$ be a hereditary category with Serre duality, generated by projective objects. Then $\mathcal{A} \cong \operatorname{rep} Q$ for a strongly locally finite thread quiver Q.*

Proof. Let $\mathfrak{a}$ be the category of projectives of $\mathcal{A}$, thus $\mathfrak{a}$ is a finite k-variety. Since $\mathfrak{a}$ generates $\mathcal{A}$, the fully faithful functor $\operatorname{mod} \mathfrak{a} \to \mathcal{A}$ is an equivalence. By definition $\mathfrak{a}$ is semi-hereditary and Theorem 4.7 yields that $\mathfrak{a}$ is a dualizing k-variety. The result now follows from Theorem 7.18. $\square$

With a thread quiver Q, there is thus an associated 2-functor $\mathfrak{a} : \operatorname{Free}(\cdot \leftarrow \cdot \to \cdot) \longrightarrow k\mathfrak{AdCat}$ given by

$$\begin{tikzcd}
\bigoplus_{t\in Q_t} k(\cdot \to \cdot) \arrow[r, "f"] \arrow[d, "g"'] & kQ_r \\
\bigoplus_{t\in Q_t} k\mathcal{L}_t &
\end{tikzcd}$$

The category $\operatorname{Rep}_k Q$ is equivalent to $\operatorname{Hom}_{2\mathcal{F}}(\mathfrak{a}, \operatorname{Mod} k)$. Thus the objects of the category $\operatorname{Rep}_k Q$ are given by

(1) an object $N(-) : kQ_r \to \operatorname{Mod} k$ of $\operatorname{Mod} kQ_r$,
(2) for every thread t, an object $L_t(-) : \mathcal{L}_t \to \operatorname{Mod} k$ of $\operatorname{Mod} k\mathcal{L}_t$, and
(3) a natural equivalence $\alpha : \oplus_t L_t(g-) \Rightarrow N(f-)$.

The morphisms are given by the modifications, thus given the data $(N, \{L_t\}_t, \alpha)$ and $(N', \{L'_t\}_t, \alpha')$ of two representations, a morphism is given by

(1) a natural transformation $\beta : N \Rightarrow N'$,
(2) a natural transformation $\gamma : \oplus_t L_t \Rightarrow \oplus_t L'_t$

such that the following diagram commutes.

$$\begin{tikzcd}
\oplus_t L_t(g-) \arrow[r, "\alpha"] \arrow[d, "\gamma\bullet 1_g"'] & N(f-) \arrow[d, "\beta\bullet 1_f"] \\
\oplus_t L'_t(g-) \arrow[r, "\alpha'"'] & N'(f-)
\end{tikzcd}$$

As in the proof of Proposition 7.16, the category $\operatorname{rep}_k Q$ has a similar description obtained by requiring N and L_t to be finitely presented representations. The category $\operatorname{rep}_k Q$ will be abelian and hereditary. When Q is a strongly locally finite quiver, then $\operatorname{rep}_k Q$ has Serre duality.

Note that even when $\operatorname{mod} kQ_r$ and $\operatorname{mod} (\oplus_{t\in Q_t} k\mathcal{L}_t)$ are well-understood, the last commutative diagram that is required can make the category $\operatorname{rep}_k Q$ contain a wild hereditary category.

Example 8.2. The category of finitely generated representations of the thread quiver

$$\begin{tikzcd}
 & & \cdot & \\
\cdot \arrow[r] & \cdot \arrow[r] & \cdot \arrow[u] \arrow[r, dotted] & \cdot
\end{tikzcd}$$

contains the representations of a wild quiver as a full and exact subcategory.

$$\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[dd, equal, bend left] \\
\mathfrak{a} \arrow[dr, bend right, "f\circ g_L"'] \arrow[ur, phantom, "\Nearrow"] & \\
 & \mathfrak{b}
\end{tikzcd}
\qquad
\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, "i"] \arrow[ddr, equal, bend left] & \\
\mathfrak{a} \arrow[r, "j"'] \arrow[drr, bend right, "f\circ g_L"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p} \arrow[dr, dashed, "i_L"] \arrow[ur, phantom, "\overset{\gamma}{\Swarrow}"] & \\
 & \arrow[u, phantom, "\overset{\beta}{\Nearrow}"] & \mathfrak{b}
\end{tikzcd}$$

FIGURE 4. Construction of i_L

APPENDIX A. 2-PUSHOUTS

Let $\mathcal{I}$ be the 1-category Free$(2 \xleftarrow{s} 1 \xrightarrow{t} 3)$ and $\mathfrak{a} : \mathcal{I} \to k\mathfrak{AdCat}$ a 2-functor. The 2-colimit of $\mathfrak{a}$ is called a *2-pushout.*

To specify a 2-natural transformation from $\mathfrak{a}$ to $\mathcal{C} \in k\mathfrak{AdCat}$, one needs three functors ($\mathfrak{f}_i : \mathfrak{a}(i) \to \mathcal{C}$, i=1,2,3) and two natural equivalences ($\mathfrak{f}_1 \overset{\sim}{\Rightarrow} \mathfrak{f}_2 \circ \mathfrak{a}(s)$ and $\mathfrak{f}_1 \overset{\sim}{\Rightarrow} \mathfrak{f}_3 \circ \mathfrak{a}(t)$). Up to invertible modification, we may assume $\mathfrak{f}_1 = \mathfrak{f}_2 \circ \mathfrak{a}(s)$ such that it suffices to give only the functors $\mathfrak{f}_2$ and $\mathfrak{f}_3$, and a natural equivalence $\mathfrak{f}_2 \circ \mathfrak{a}(s) \overset{\sim}{\Rightarrow} \mathfrak{f}_3 \circ \mathfrak{a}(t)$.

The following theorem shows a connection between 2-pushouts and adjoints.

Theorem A.1. *Consider the 2-pushout*

$$\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, dashed, "i"] \\
\mathfrak{c} \arrow[r, dashed, "j"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p}
\end{tikzcd}$$

in any 2-category. If g is fully faithful and has a left (right) adjoint, then the same holds for i.

Proof. Consider the first diagram in Figure 4, where the natural transformation is given by

$$1_f \bullet \epsilon_g : f \circ g_L \circ g \overset{\sim}{\Rightarrow} f,$$

and where $\epsilon_g : g_L \circ g \Rightarrow 1$ is the counit of the adjunction (g_L, g). We can complete this diagram as shown in Figure 4. This gives the following identity

$$(\gamma^{-1} \bullet 1_f) \circ (1_{i_L} \bullet \alpha) \circ (\beta \bullet 1_g) = 1_f \bullet \epsilon_g. \tag{4}$$

We claim $i_L : \mathfrak{p} \to \mathfrak{b}$ is a left adjoint to i. To prove this, we shall define a unit $\eta : 1_{\mathfrak{p}} \Rightarrow i \circ i_L$ and a counit $\epsilon : i_L \circ i \Rightarrow 1_{\mathfrak{b}}$ and show that

$$(1 \bullet \epsilon) \circ (\eta \bullet 1) = 1 \text{ and } (\epsilon \bullet 1) \circ (1 \bullet \eta) = 1.$$

We define $\epsilon = \gamma^{-1} : i_L \circ i \overset{\sim}{\Rightarrow} 1_{\mathfrak{b}}$ with γ as in the second diagram in Figure 4. For the definition of η, consider the diagrams in Figure 5, where the natural transformations on the right-hand side are given by

$$\begin{aligned}
1_i \bullet \gamma \quad &: \quad i \Rightarrow i \circ i_L \circ i \\
\delta = (1_i \bullet \beta) \circ (\alpha \bullet 1_{g_L}) \quad &: \quad j \circ g \circ g_L \Rightarrow i \circ f \circ g_L \Rightarrow i \circ i_L \circ j
\end{aligned}$$

There is a modification, going from the 2-natural transformation on the left-hand side of Figure 5 to the one on the right-hand side, given by

$$\begin{aligned}
1_i \quad &: \quad i \Rightarrow i \\
1_j \bullet \eta_g \quad &: \quad j \Rightarrow j \circ g \circ g_L
\end{aligned}$$

$$\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, dashed, "i"] \arrow[ddr, bend left, "i"] & \\
\mathfrak{a} \arrow[r, dashed, "j"'] \arrow[drr, bend right, "j"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p} \arrow[dr, equal] \arrow[r, phantom, "="] \arrow[d, phantom, "="] & \\
& & \mathfrak{p}
\end{tikzcd}
\qquad
\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, dashed, "i"] \arrow[ddr, bend left, "i"] & \\
\mathfrak{a} \arrow[r, dashed, "j"'] \arrow[drr, bend right, "j\circ g\circ g_L"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p} \arrow[dr, dashed, "i\circ i_L"] \arrow[r, phantom, "\underset{1\bullet\gamma}{\Swarrow}"] \arrow[d, phantom, "\overset{\delta}{\Nearrow}"] & \\
& & \mathfrak{p}
\end{tikzcd}$$

Figure 5. Construction of unit

$$\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, dashed, "i"] \arrow[dddrr, bend left, "i_L\circ i"] & & \\
\mathfrak{c} \arrow[r, dashed, "j"'] \arrow[ddrrr, bend right, "i_L\circ j"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p} \arrow[dr, equal] \arrow[r, phantom, "="] \arrow[d, phantom, "="] & & \\
& & \mathfrak{p} \arrow[dr, "i_L"] & \\
& & & \mathfrak{b}
\end{tikzcd}
\qquad
\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, dashed, "i"] \arrow[dddrr, bend left, "i_L\circ i"] & & \\
\mathfrak{c} \arrow[r, dashed, "j"'] \arrow[ddrrr, bend right, "i_L\circ j\circ g\circ g_L"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p} \arrow[dr, "i\circ i_L"] \arrow[r, phantom, "\Swarrow"] \arrow[d, phantom, "\Nearrow"] & & \\
& & \mathfrak{p} \arrow[dr, "i_L"] & \\
& & & \mathfrak{b}
\end{tikzcd}$$

Figure 6. Diagrams occurring in the definition of the modification Λ

where $\eta_g : 1 \Rightarrow g \circ g_L$ is the unit of the adjunction (g_L, g). This modification corresponds to a natural transformation $\eta : 1_{\mathfrak{p}} \Rightarrow i \circ i_L$, satisfying the following commutative diagrams (see Remark 2.8)

$$\begin{tikzcd}
i \arrow[r, equal] \arrow[d, equal] & i \arrow[d, Rightarrow, "1\bullet\gamma"] \\
i \arrow[r, Rightarrow, "\eta\bullet 1"'] & i \circ i_L \circ i
\end{tikzcd}
\qquad
\begin{tikzcd}
j \arrow[r, Rightarrow, "1\bullet\eta_g"] \arrow[d, equal] & j \circ g \circ g_L \arrow[d, Rightarrow, "\delta"] \\
j \arrow[r, Rightarrow, "\eta\bullet 1_j"'] & i \circ i_L \circ j
\end{tikzcd}$$

From the first commutative diagram, we obtain

$$(1 \bullet \epsilon) \circ (\eta \bullet 1) = (1 \bullet \gamma)^{-1} \circ (\eta \bullet 1) = 1.$$

To show the other equation, and establish the adjointness of i and i_L, we add the functor $i_L : \mathfrak{p} \to \mathfrak{b}$ to the diagrams in Figure 5 (see Figure 6). The natural transformations of the second diagram of Figure 6 are given by

$$\begin{aligned}
1_{i_L} \bullet \delta &: i_L \circ j \circ g \circ g_L \Rightarrow i_L \circ i \circ i_L \circ j \\
1_{i_L} \bullet 1_i \bullet \gamma &: i_L \circ i \Rightarrow i_L \circ i \circ i_L \circ i
\end{aligned}$$

$$\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, dashed, "i"] \arrow[ddrr, bend left, "i_L \circ i"] & & \\
\mathfrak{c} \arrow[r, dashed, "j"'] \arrow[drrr, bend right, "i_L \circ j \circ g \circ g_L"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p} \arrow[dr, "i_L"] & \Swarrow & \\
& \Nearrow & \mathfrak{b} \arrow[dr, "i_L \circ i"] & \\
& & & \mathfrak{b}
\end{tikzcd}
\qquad
\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] & \mathfrak{b} \arrow[d, dashed, "i"] \arrow[ddrr, bend left, "i_L \circ i"] & & \\
\mathfrak{c} \arrow[r, dashed, "j"'] \arrow[drrr, bend right, "i_L \circ j"'] \arrow[ur, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{p} \arrow[dr, "i_L"] & /\!/ & \\
& /\!/ & \mathfrak{b} \arrow[dr, equal] & \\
& & & \mathfrak{b}
\end{tikzcd}$$

FIGURE 7. Diagrams occurring in the definition of the modification Ω

The modification going from the diagram on the left to the diagram on the right in Figure 5 induces a modification Λ given by

$$\begin{array}{rcl} 1_{i_L} \bullet 1_j \bullet \eta_g & : & i_L \circ j \Rightarrow i_L \circ j \circ g \circ g_L \\ 1 & : & i_L \circ i \Rightarrow i_L \circ i \end{array}$$

which corresponds to the natural transformation $1 \bullet \eta : i_L \Rightarrow i_L \circ i \circ i_L$.

Likewise, there is a modification Ω (going from the 2-natural transformation on the left hand side of Figure 7 to the 2-natural transformation on the right hand side) given by

$$\begin{array}{rcl} (\epsilon \bullet 1) \circ (1 \bullet \delta) & : & i_L \circ j \circ g \circ g_L \Rightarrow i_L \circ j \\ 1 & : & i_L \circ i \Rightarrow i_L \circ i \end{array}$$

which induces the following commutative diagrams

$$\begin{tikzcd}
i_L \circ i \arrow[r, Rightarrow, "\gamma \bullet 1_{i_L} \bullet 1_i"] \arrow[d, equal] & i_L \circ i \circ i_L \circ i \arrow[d, Rightarrow, "\epsilon \bullet 1_{i_L} \bullet 1_i"] \\
i_L \circ i \arrow[r, equal] & i \circ i_L
\end{tikzcd}
\qquad
\begin{tikzcd}
i_L \circ j \circ g \circ g_L \arrow[r, Rightarrow, "1_{i_L} \bullet \delta"] \arrow[d, Rightarrow, "(\epsilon \bullet 1_{i_L} \bullet 1_j) \circ (1_{i_L} \bullet \delta)"] & i_L \circ i \circ i_L \circ j \arrow[d, Rightarrow, "\epsilon \bullet 1_{i_L} \bullet 1_j"] \\
i_L \circ j \arrow[r, equal] & i_L \circ j
\end{tikzcd}$$

such that Ω corresponds to the natural transformation $\epsilon \bullet 1_{i_L} : i_L \circ i \circ i_L \to i_L$.

We claim that $\Omega \circ \Lambda = 1$. The only nontrivial fact to check is that $\Omega \circ \Lambda$ induces the identity on $i_L \circ j : \bigoplus_t k\mathcal{L}_t \to \mathfrak{p}$. This follows from

$$\begin{array}{rl} & (\epsilon \bullet 1_{i_L} \bullet 1_j) \circ (1_{i_L} \bullet \delta) \circ (1_{i_L} \bullet 1_j \bullet \eta_g) \\ = & (\epsilon \bullet 1_{i_L} \bullet 1_j) \circ (1_{i_L} \bullet 1_i \bullet \beta) \circ (1_{i_L} \bullet \alpha \bullet 1_{g_L}) \circ (1_{i_L} \bullet 1_j \bullet \eta_g) \\ = & \beta \circ (\epsilon \bullet 1_f \bullet 1_{g_L}) \circ (1_{i_L} \bullet \alpha \bullet 1_{g_L}) \circ (1_{i_L} \bullet 1_j \bullet \eta_g) \\ = & \beta \circ (1_f \bullet \epsilon_g \bullet 1_{g_L}) \circ (\beta^{-1} \bullet 1_g \bullet 1_{g_L}) \circ (1_{i_L} \bullet 1_j \bullet \eta_g) \\ = & \beta \circ (1_f \bullet \epsilon_g \bullet 1_{g_L}) \circ (1_f \bullet 1_{g_L} \bullet \eta_g) \circ \beta^{-1} \\ = & \beta \circ (1_f \bullet 1_{g_L}) \circ \beta^{-1} \\ = & 1_{i_L} \bullet 1_j \end{array}$$

where we have used equations (1) and (4). Since $\Omega \circ \Lambda = 1$, we find that $(\epsilon \bullet 1) \circ (1 \bullet \eta) = 1$ and hence we may conclude that i_L is indeed left adjoint to i. Since ϵ is invertible, the functor i is fully faithful.

$$\begin{tikzcd}
\mathfrak{a} \arrow[r, "f"] \arrow[d, "g"'] \arrow[dr, phantom, "\overset{\alpha}{\Nearrow}"] & \mathfrak{b} \arrow[d, "i"] \arrow[ddr, equal, bend left] \arrow[ddr, phantom, "\Swarrow"] & \\
\mathfrak{c} \arrow[r, "j"'] \arrow[drr, bend right, "j \circ g_R"'] \arrow[drr, phantom, "\Nearrow"] & \mathfrak{p} \arrow[dr, dashed, "i_R"] & \\
& & \mathfrak{b}
\end{tikzcd}$$

Figure 8. Construction of i_R

Similarly, one verifies that the functor i_R, defined in Figure 8, is right adjoint to i. □

References

1. Maurice Auslander, *Representation theory of Artin algebras. I*, Comm. Algebra **1** (1974), 177–268.
2. Maurice Auslander and Idun Reiten, *Stable equivalence of dualizing R-varieties*, Advances in Math. **12** (1974), 306–366.
3. ———, *Stable equivalence of dualizing R-varieties. II. Hereditary dualizing R-varieties*, Advances in Math. **17** (1975), 93–121.
4. Maurice Auslander and Sverre O. Smalø, *Almost split sequences in subcategories*, J. Algebra **69** (1981), no. 2, 426–454.
5. Carl Fredrik Berg and Adam-Christiaan van Roosmalen, *Hereditary categories with Serre duality which are generated by preprojectives.*
6. ———, *The quiver of projectives in hereditary categories with Serre duality*, J. Pure Appl. Algebra, accepted.
7. A. I. Bondal and M. M. Kapranov, *Representable functors, Serre functors, and reconstructions*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1183–1205, 1337.
8. Francis Borceux, *Handbook of categorical algebra. 1*, Encyclopedia of Mathematics and its Applications, vol. 50, Cambridge University Press, Cambridge, 1994, Basic category theory.
9. ———, *Handbook of categorical algebra. 2*, Encyclopedia of Mathematics and its Applications, vol. 51, Cambridge University Press, Cambridge, 1994, Categories and structures.
10. Xiao-Wu Chen, *Generalized Serre duality*, preprint.
11. Dieter Happel, *Triangulated categories in the representation theory of finite dimensional algebras*, London Mathematical Society Lecture Notes Series, vol. 119, Cambridge University Press, Cambridge, 1988.
12. Osamu Iyama and Yuji Yoshino, *Mutations in triangulated categories and rigid Cohen-Macaulay modules*, Invent. Math **172** (2008), no. 1, 117–168.
13. G.M. Kelly, *Basic concepts of enriched category theory*, Repr. Theory Appl. Categ. (2005), no. 10, vi+137 pp. (electronic), Reprint of the 1982 original [Cambridge Univ. Press, Cambridge].
14. Saunders MacLane, *Categories for the working mathematician*, Graduate Texts in Mathematics, Vol. 5, Springer-Verlag, New York, 1971.
15. Claus Michael Ringel, *The diamond category of a locally discrete ordered set*, Representations of algebra. Vol. I, II, Beijing Norm. Univ. Press, Beijing, 2002, pp. 387–395.
16. Sverre O. Smalø, *Almost split sequences in categories of representations of quivers II*, Representations of algebras (São Paulo, 1999), Lecture Notes in Pure and Appl. Math.
17. Adam-Christiaan van Roosmalen, *Classification of abelian hereditary directed categories satisfying Serre duality*, Trans. Amer. Math. Soc. **360** (2008), no. 5, 2467–2503.
18. Ingo Waschkies, *The stack of microlocal perverse sheaves*, Bull. Soc. Math. France **132** (2004), no. 3, 397–462.

Carl Fredrik Berg, Institutt for matematiske fag, NTNU, 7491 Trondheim, Norway, (Currently working for StatoilHydro R&D Centre, Arkitekt Ebbells veg 10, Rotvoll, 7053 Ranheim, Norway)
E-mail address: `carlpaatur@hotmail.com`

Adam-Christiaan van Roosmalen, Mathematisches Institut, Universität Bonn, Endenicher Allee 60, 53111 Bonn, Germany
E-mail address: `vroosmal@math.uni-bonn.de`